

Optimal Stopping for Dynamic Convex Risk Measures

Erhan Bayraktar*, Ioannis Karatzas†, Song Yao‡

Abstract

We use martingale and stochastic analysis techniques to study a continuous-time optimal stopping problem, in which the decision maker uses a dynamic convex risk measure to evaluate future rewards. We also find a saddle point for an equivalent zero-sum game of control and stopping, between an agent (the “stopper”) who chooses the termination time of the game, and an agent (the “controller”, or “nature”) who selects the probability measure.

Keywords: Convex risk measures, continuous-time optimal stopping, robustness methods, zero sum games, saddle point, reflected backward stochastic differential equations, BMO martingales.

1 Introduction

Let us consider a complete, filtered probability space $(\Omega, \mathcal{F}, P)$, $\mathbf{F} = \{\mathcal{F}_t\}_{t \geq 0}$, and on it a bounded, adapted process Y that satisfies certain regularity conditions. Given any stopping time ν of the filtration $\mathbf{F}$, our goal is to find a stopping time $\tau_*(\nu) \in \mathcal{S}_{\nu, T}$ that satisfies

$$\operatorname{ess\,inf}_{\gamma \in \mathcal{S}_{\nu, T}} \rho_{\nu, \gamma}(Y_\gamma) = \rho_{\nu, \tau_*(\nu)}(Y_{\tau_*(\nu)}), \quad P - \text{a.s.} \quad (1.1)$$

Here $\mathcal{S}_{\nu, T}$ is the set of stopping times γ satisfying $\nu \leq \gamma \leq T$, P -a.s., and the collection of functionals $\{\rho_{\nu, \gamma} : \mathbb{L}^\infty(\mathcal{F}_\gamma) \rightarrow \mathbb{L}^\infty(\mathcal{F}_\nu)\}_{\nu \in \mathcal{S}_{0, T}, \gamma \in \mathcal{S}_{\nu, T}}$ is a “dynamic convex risk measure” in the sense of Delbaen et al. [2009]. Our motivation is to solve the optimal stopping problem of a decision maker who evaluates future rewards/risks using dynamic convex risk measures rather than statistical expectations. This question can also be cast as a *robust optimal stopping* problem, in which the decision maker has to act in the presence of so-called “Knightian uncertainty” regarding the underlying probability measure.

When the filtration $\mathbf{F}$ is generated by a Brownian motion, the dynamic convex risk measure admits the following representation: There exists a suitable nonnegative function f , convex in its spatial argument, such that the representation

$$\rho_{\nu, \gamma}(\xi) = \operatorname{ess\,sup}_{Q \in \mathcal{Q}_\nu} E_Q \left[-\xi - \int_\nu^\gamma f(s, \theta_s^Q) ds \mid \mathcal{F}_\nu \right], \quad P - \text{a.s.}$$

holds for all $\xi \in \mathbb{L}^\infty(\mathcal{F}_\gamma)$. Here $\mathcal{Q}_\nu$ is the collection of probability measures Q which are equivalent to P on $\mathcal{F}$, equal to P on $\mathcal{F}_\nu$, and satisfy a certain integrability condition; whereas θ^Q is the predictable process whose stochastic exponential gives the density of Q with respect to P . In this setting we establish a minimax result, namely

$$V(\nu) \triangleq \operatorname{ess\,sup}_{\gamma \in \mathcal{S}_{\nu, T}} \left(\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \mid \mathcal{F}_\nu \right] \right) = \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} \left(\operatorname{ess\,sup}_{\gamma \in \mathcal{S}_{\nu, T}} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \mid \mathcal{F}_\nu \right] \right), \quad (1.2)$$

*Department of Mathematics, University of Michigan, Ann Arbor, MI 48109; email: erhan@umich.edu. This author is supported in part by the National Science Foundation, under grant number DMS-0906257.

†INTECH Investment Management, One Palmer Square, Suite 441, Princeton, NJ 08542; e-mail: ik@enhanced.com; and Department of Mathematics, Columbia University, New York, NY 10027; e-mail: ik@math.columbia.edu. Research supported in part by the National Science Foundation under Grant NSF-DMS-09-05754.

‡Department of Mathematics, University of Michigan, Ann Arbor, MI 48109; email: songyao@umich.edu.

and construct an optimal stopping time $\tau(\nu)$ as the limit of stopping times that are optimal under expectation criteria — see Theorem 3.1. We show that the process $\{\mathbf{1}_{\{t \geq \nu\}} V(\tau(\nu) \wedge t)\}_{t \in [0, T]}$ admits an RCLL modification $V^{0, \nu}$ such that for any $\gamma \in \mathcal{S}_{0, T}$, we have $V_\gamma^{0, \nu} = \mathbf{1}_{\{\gamma \geq \nu\}} V(\tau(\nu) \wedge \gamma)$, P -a.s. We show that the stopping time $\tau_V(\nu) \triangleq \inf \{t \in [\nu, T] : V_t^{0, \nu} = Y_t\}$ attains the infimum in (1.1). Finally, we construct a saddle point of the stochastic game in (1.2).

The discrete-time optimal stopping problem for coherent risk measures was studied by Föllmer and Schied [2004, Section 6.5] and Cheridito et al. [2006, Sections 5.2 and 5.3]. Delbaen [2006] and Karatzas and Zamfirescu [2006], on the other hand, considered continuous-time optimal stopping problems in which the essential infimum over the stopping times in (1.1) is replaced by an essential supremum. The controller-and-stopper problem of Lepeltier [1985] and Karatzas and Zamfirescu [2008], and the optimal stopping for non-linear expectations in Bayraktar and Yao [2009], are the closest in spirit to our work. However, since our assumptions on the random function f and the set $\mathcal{Q}_\nu$ are dictated by the representation theorem for dynamic convex risk measures, the results in these papers cannot be directly applied. In particular, because of the integrability assumption that appears in the definition of $\mathcal{Q}_\nu$ (subsection 1.1), this set may not be closed under *pasting*; see Remark 3.3. Moreover, the extant results on controller-and-stopper games would require that f and the θ^Q 's be bounded. We overcome these technical difficulties by using approximation arguments which rely on *truncation* and *localization* techniques. On the other hand, in finding a saddle point Karatzas and Zamfirescu [2008] used the weak compactness of the collection of probability measures, in particular the boundedness of θ^Q 's. We avoid making this assumption by using techniques from Reflected Backward Stochastic Differential Equations (RBSDEs). In particular, using a comparison theorem and the fact that V can be approximated by solutions of BSDEs with Lipschitz generators, we show that V solves a quadratic RBSDE (QRBSDE). The relationship between the solutions of QRBSDEs and the BMO martingales helps us construct a saddle point. We should point out that the convexity of f is not needed to derive our results; cf. Remark 3.1.

The layout of the paper is simple. In Section 2 we recall the definition of the dynamic convex risk measures and a representation theorem. We solve the optimal stopping problem in Section 3. In Section 4 we find a saddle point for the stochastic controller-and-stopper game in (1.2). The proofs of our results are given in Section 5.

1.1 Notation and Preliminaries

Throughout this paper we let B be a d -dimensional Brownian Motion defined on the probability space $(\Omega, \mathcal{F}, P)$, and consider the augmented filtration generated by it, i.e.,

$$\mathbf{F} = \{\mathcal{F}_t \triangleq \sigma(B_s; s \in [0, t]) \vee \mathcal{N}\}_{t \geq 0}, \text{ where } \mathcal{N} \text{ is the collection of all } P\text{-null sets in } \mathcal{F}.$$

We fix a finite time horizon $T > 0$, denote by $\mathcal{P}$ (resp. $\widehat{\mathcal{P}}$) the predictably (resp. progressively) measurable σ -field on $\Omega \times [0, T]$, and let $\mathcal{S}_{0, T}$ be the set of all $\mathbf{F}$ -stopping times ν such that $0 \leq \nu \leq T$, P -a.s. From now on, when writing $\nu \leq \gamma$, we always mean two stopping times $\nu, \gamma \in \mathcal{S}_{0, T}$ such that $\nu \leq \gamma$, P -a.s. For any $\nu \leq \gamma$ we define $\mathcal{S}_{\nu, \gamma} \triangleq \{\sigma \in \mathcal{S}_{0, T} \mid \nu \leq \sigma \leq \gamma, P\text{-a.s.}\}$ and let $\mathcal{S}_{\nu, \gamma}^*$ denote all finite-valued stopping times in $\mathcal{S}_{\nu, \gamma}$.

The following spaces of functions will be used in the sequel:

- Let $\mathcal{G}$ be a generic sub- σ -field of $\mathcal{F}$. $\mathbb{L}^0(\mathcal{G})$ denotes the space of all real-valued, $\mathcal{G}$ -measurable random variables.
- $\mathbb{L}^\infty(\mathcal{G}) \triangleq \{\xi \in \mathbb{L}^0(\mathcal{G}) : \|\xi\|_\infty \triangleq \operatorname{ess\,sup}_{\omega \in \Omega} |\xi(\omega)| < \infty\}$.
- $\mathbb{L}_{\mathbf{F}}^0[0, T]$ denotes the space of all real-valued, $\mathbf{F}$ -adapted processes.
- $\mathbb{L}_{\mathbf{F}}^\infty[0, T] \triangleq \{X \in \mathbb{L}_{\mathbf{F}}^0[0, T] : \|X\|_\infty \triangleq \operatorname{ess\,sup}_{(t, \omega) \in [0, T] \times \Omega} |X_t(\omega)| < \infty\}$.
- $\mathbb{C}_{\mathbf{F}}^p[0, T] \triangleq \{X \in \mathbb{L}_{\mathbf{F}}^p[0, T] : X \text{ has continuous paths}\}$, $p = 0, \infty$.
- $\mathbb{C}_{\mathbf{F}}^2[0, T] \triangleq \{X \in \mathbb{C}_{\mathbf{F}}^0[0, T] : E\left(\sup_{t \in [0, T]} |X_t|^2\right) < \infty\}$.

- $\mathbb{H}_{\mathbf{F}}^2([0, T]; \mathbb{R}^d)$ (resp. $\widehat{\mathbb{H}}_{\mathbf{F}}^2([0, T]; \mathbb{R}^d)$) denotes the space of all $\mathbb{R}^d$ -valued, $\mathbf{F}$ -adapted predictably (resp. progressively) measurable processes X with $E \int_0^T |X_t|^2 dt < \infty$.
- $\mathbb{H}_{\mathbf{F}}^\infty([0, T]; \mathbb{R}^d)$ denotes the space of all $\mathbb{R}^d$ -valued, $\mathbf{F}$ -adapted predictably measurable processes X with $\operatorname{ess\,sup}_{(t, \omega) \in [0, T] \times \Omega} |X_t(\omega)| < \infty$.
- $\mathbb{K}_{\mathbf{F}}[0, T]$ denotes the space of all real-valued, $\mathbf{F}$ -adapted continuous increasing processes K with $K_0 = 0$.

Let us consider the set $\mathcal{M}^e$ of all probability measures on $(\Omega, \mathcal{F})$ which are equivalent to P . For any $Q \in \mathcal{M}^e$, it is well-known that there is an $\mathbb{R}^d$ -valued predictable process θ^Q with $\int_0^T |\theta_t^Q|^2 dt < \infty$, P -a.s., such that the density process Z^Q of Q with respect to P is the stochastic exponential of θ^Q , namely,

$$Z_t^Q = \mathcal{E}(\theta^Q \bullet B)_t = \exp \left\{ \int_0^t \theta_s^Q dB_s - \frac{1}{2} \int_0^t |\theta_s^Q|^2 ds \right\}, \quad 0 \leq t \leq T.$$

We denote $Z_{\nu, \gamma}^Q \triangleq Z_\gamma^Q / Z_\nu^Q = \exp \left\{ \int_\nu^\gamma \theta_s^Q dB_s - \frac{1}{2} \int_\nu^\gamma |\theta_s^Q|^2 ds \right\}$ for any $\nu \leq \gamma$. Moreover, for any $\nu \in \mathcal{S}_{0, T}$ and with the notation $\llbracket 0, \nu \rrbracket \triangleq \{(t, \omega) \in [0, T] \times \Omega : 0 \leq t < \nu(\omega)\}$ for the stochastic interval, we define

$$\begin{aligned} \mathcal{P}_\nu &\triangleq \{Q \in \mathcal{M}^e : Q = P \text{ on } \mathcal{F}_\nu\} = \{Q \in \mathcal{M}^e : \theta_t^Q(\omega) = 0, \text{ dt} \otimes dP\text{-a.e. on } \llbracket 0, \nu \rrbracket\}, \\ \mathcal{Q}_\nu &\triangleq \{Q \in \mathcal{P}_\nu : E_Q \int_\nu^T f(s, \theta_s^Q) ds < \infty\}. \end{aligned}$$

2 Dynamic Convex Risk Measures

Definition 2.1. A dynamic convex risk measure is a family of functionals $\{\rho_{\nu, \gamma} : \mathbb{L}^\infty(\mathcal{F}_\gamma) \rightarrow \mathbb{L}^\infty(\mathcal{F}_\nu)\}_{\nu \leq \gamma}$ which satisfy the following properties: For any stopping times $\nu \leq \gamma$ and any $\mathbb{L}^\infty(\mathcal{F}_\gamma)$ -measurable random variables ξ, η , we have

- “Monotonicity”: $\rho_{\nu, \gamma}(\xi) \leq \rho_{\nu, \gamma}(\eta)$, P -a.s. if $\xi \geq \eta$, P -a.s.
- “Translation Invariance”: $\rho_{\nu, \gamma}(\xi + \eta) = \rho_{\nu, \gamma}(\xi) - \eta$, P -a.s. if $\eta \in \mathbb{L}^\infty(\mathcal{F}_\nu)$.
- “Convexity”: $\rho_{\nu, \gamma}(\lambda \xi + (1 - \lambda)\eta) \leq \lambda \rho_{\nu, \gamma}(\xi) + (1 - \lambda) \rho_{\nu, \gamma}(\eta)$, P -a.s. for any $\lambda \in (0, 1)$.
- “Normalization”: $\rho_{\nu, \gamma}(0) = 0$, P -a.s.

Delbaen et al. [2009] provide a representation result, Proposition 2.1 below, for dynamic convex risk measures $\{\rho_{\nu, \gamma}\}_{\nu \leq \gamma}$ that satisfy the following properties:

- (A1) “Continuity from above”: For any decreasing sequence $\{\xi_n\} \subset \mathbb{L}^\infty(\mathcal{F}_\gamma)$ with $\xi \triangleq \lim_{n \rightarrow \infty} \downarrow \xi_n \in \mathbb{L}^\infty(\mathcal{F}_\gamma)$, it holds P -a.s. that $\lim_{n \rightarrow \infty} \uparrow \rho_{\nu, \gamma}(\xi_n) = \rho_{\nu, \gamma}(\xi)$.
- (A2) “Time Consistency”: For any $\sigma \in \mathcal{S}_{\nu, \gamma}$ we have: $\rho_{\nu, \sigma}(-\rho_{\sigma, \gamma}(\xi)) = \rho_{\nu, \gamma}(\xi)$, P -a.s.
- (A3) “Zero-One Law”: For any $A \in \mathcal{F}_\nu$, we have: $\rho_{\nu, \gamma}(\mathbf{1}_A \xi) = \mathbf{1}_A \rho_{\nu, \gamma}(\xi)$, P -a.s.
- (A4) $\operatorname{ess\,inf}_{\xi \in \mathcal{A}_t} E_P[\xi | \mathcal{F}_t] = 0$, where $\mathcal{A}_t \triangleq \{\xi \in \mathbb{L}^\infty(\mathcal{F}_T) : \rho_{t, T}(\xi) \leq 0\}$.

Proposition 2.1. Let $\{\rho_{\nu, \gamma}\}_{\nu \leq \gamma}$ be a dynamic convex risk measure satisfying (A1)-(A4). Then for any $\nu \leq \gamma$ and $\xi \in \mathbb{L}^\infty(\mathcal{F}_\gamma)$, we have

$$\rho_{\nu, \gamma}(\xi) = \operatorname{ess\,sup}_{Q \in \mathcal{Q}_\nu} E_Q \left[-\xi - \int_\nu^\gamma f(s, \theta_s^Q) ds \mid \mathcal{F}_\nu \right], \quad P\text{-a.s.} \quad (2.1)$$

Here $f : [0, T] \times \Omega \times \mathbb{R}^d \rightarrow [0, \infty]$ is a suitable measurable function, such that

- (f1) $f(\cdot, \cdot, z)$ is predictable for any $z \in \mathbb{R}^d$;
- (f2) $f(t, \omega, \cdot)$ is proper convex, and lower semi-continuous for $dt \otimes dP$ -a.e. $(t, \omega) \in [0, T] \times \Omega$; and
- (f3) $f(t, \omega, 0) = 0$, $dt \otimes dP$ -a.e.

We refer to Rockafellar [1997], p. 24 for the notion of “proper convex function”, and review some basic properties of the essential extrema as in Neveu [1975, Proposition VI-1-1] or Föllmer and Schied [2004, Theorem A.32].

Lemma 2.1. *Let $\{\xi_i\}_{i \in \mathcal{I}}$ and $\{\eta_i\}_{i \in \mathcal{I}}$ be two classes of $\mathcal{F}$ -measurable random variables with the same index set $\mathcal{I}$.*

- (1) *If $\xi_i \leq (=) \eta_i$, P -a.s. holds for all $i \in \mathcal{I}$, then $\operatorname{ess\,sup}_{i \in \mathcal{I}} \xi_i \leq (=) \operatorname{ess\,sup}_{i \in \mathcal{I}} \eta_i$, P -a.s.*
- (2) *For any $A \in \mathcal{F}$, it holds P -a.s. that $\operatorname{ess\,sup}_{i \in \mathcal{I}} (\mathbf{1}_A \xi_i + \mathbf{1}_{A^c} \eta_i) = \mathbf{1}_A \operatorname{ess\,sup}_{i \in \mathcal{I}} \xi_i + \mathbf{1}_{A^c} \operatorname{ess\,sup}_{i \in \mathcal{I}} \eta_i$. In particular, $\operatorname{ess\,sup}_{i \in \mathcal{I}} (\mathbf{1}_A \xi_i) = \mathbf{1}_A \operatorname{ess\,sup}_{i \in \mathcal{I}} \xi_i$, P -a.s.*
- (3) *For any $\mathcal{F}$ -measurable random variable γ and any $\lambda > 0$, we have $\operatorname{ess\,sup}_{i \in \mathcal{I}} (\lambda \xi_i + \gamma) = \lambda \operatorname{ess\,sup}_{i \in \mathcal{I}} \xi_i + \gamma$, P -a.s.*

Moreover, (1)-(3) hold when we replace $\operatorname{ess\,sup}_{i \in \mathcal{I}}$ by $\operatorname{ess\,inf}_{i \in \mathcal{I}}$.

3 The Optimal Stopping Problem

In this section we study the optimal stopping problem for dynamic convex risk measures. More precisely, given $\nu \in \mathcal{S}_{0,T}$, we seek an optimal stopping time $\tau_*(\nu) \in \mathcal{S}_{\nu,T}$ that satisfies (1.1). We shall assume throughout that the reward process $Y \in \mathbb{L}_{\mathbf{F}}^\infty[0, T]$ is right-continuous and $\mathcal{Q}_0$ -quasi-left-continuous: to wit, for any increasing sequence $\{\nu_n\}_{n \in \mathbb{N}}$ in $\mathcal{S}_{0,T}$ with $\nu \stackrel{\Delta}{=} \lim_{n \rightarrow \infty} \uparrow \nu_n \in \mathcal{S}_{0,T}$, and any $Q \in \mathcal{Q}_0$, we have

$$\lim_{n \rightarrow \infty} E_Q[Y_{\nu_n} | \mathcal{F}_{\nu_1}] \leq E_Q[Y_\nu | \mathcal{F}_{\nu_1}], \quad P\text{-a.s.}$$

In light of the representation (2.1), we can alternatively express (1.1) as a *robust optimal stopping problem*, in the following sense:

$$\operatorname{ess\,sup}_{\gamma \in \mathcal{S}_{\nu,T}} \left(\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \mid \mathcal{F}_\nu \right] \right) = \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_{\tau_*(\nu)} + \int_\nu^{\tau_*(\nu)} f(s, \theta_s^Q) ds \mid \mathcal{F}_\nu \right]. \quad (3.1)$$

Remark 3.1. *We will study the robust optimal stopping problem (3.1) in a setting more general than alluded to heretofore: From now on, we only assume that $f : [0, T] \times \Omega \times \mathbb{R}^d \rightarrow [0, \infty]$ is a $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d) / \mathcal{B}([0, \infty])$ -measurable function which satisfies (f3); i.e., the convexity (f2) is not necessary for solving (3.1).*

In order to find a stopping time which is optimal, i.e., attains the essential supremum in (3.1), we introduce the lower- and upper-value, respectively, of the stochastic game suggested by (3.1), to wit, for every $\nu \in \mathcal{S}_{0,T}$:

$$\underline{V}(\nu) \stackrel{\Delta}{=} \operatorname{ess\,sup}_{\gamma \in \mathcal{S}_{\nu,T}} \left(\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \mid \mathcal{F}_\nu \right] \right), \quad \overline{V}(\nu) \stackrel{\Delta}{=} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} \left(\operatorname{ess\,sup}_{\gamma \in \mathcal{S}_{\nu,T}} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \mid \mathcal{F}_\nu \right] \right).$$

In Theorem 3.1 we shall show that the quantities $\underline{V}(\nu)$ and $\overline{V}(\nu)$ coincide at any $\nu \in \mathcal{S}_{0,T}$, i.e., a min-max theorem holds; we shall also identify two optimal stopping times in Theorems 3.1 and 3.2, respectively.

Given any probability measure $Q \in \mathcal{Q}_0$, let us introduce for each fixed $\nu \in \mathcal{S}_{0,T}$ the quantity

$$R^Q(\nu) \stackrel{\Delta}{=} \operatorname{ess\,sup}_{\zeta \in \mathcal{S}_{\nu,T}} E_Q \left[Y_\zeta + \int_\nu^\zeta f(s, \theta_s^Q) ds \mid \mathcal{F}_\nu \right] = \operatorname{ess\,sup}_{\sigma \in \mathcal{S}_{0,T}} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \mid \mathcal{F}_\nu \right] \geq Y_\nu \quad (3.2)$$

and recall from the classical theory of optimal stopping (see El Karoui [1981] or Karatzas and Shreve [1998, Appendix D]) the following result.

Proposition 3.1. *Fix a probability measure $Q \in \mathcal{Q}_0$.*

- (1) *The process $\{R^Q(t)\}_{t \in [0, T]}$ admits an RCLL modification $R^{Q,0}$ such that, for any $\nu \in \mathcal{S}_{0,T}$, we have*

$$R_\nu^{Q,0} = R^Q(\nu), \quad P\text{-a.s.} \quad (3.3)$$

(2) For every $\nu \in \mathcal{S}_{0,T}$, the stopping time $\tau^Q(\nu) \triangleq \inf\{t \in [\nu, T] : R_t^{Q,0} = Y_t\} \in \mathcal{S}_{\nu,T}$ satisfies for any $\gamma \in \mathcal{S}_{\nu, \tau^Q(\nu)}$:

$$\begin{aligned} R^Q(\nu) &= E_Q \left[Y_{\tau^Q(\nu)} + \int_{\nu}^{\tau^Q(\nu)} f(s, \theta_s^Q) ds \middle| \mathcal{F}_{\nu} \right] = E_Q \left[R^Q(\tau^Q(\nu)) + \int_{\nu}^{\tau^Q(\nu)} f(s, \theta_s^Q) ds \middle| \mathcal{F}_{\nu} \right] \\ &= E_Q \left[R^Q(\gamma) + \int_{\nu}^{\gamma} f(s, \theta_s^Q) ds \middle| \mathcal{F}_{\nu} \right], \quad P\text{-a.s.} \end{aligned} \quad (3.4)$$

Therefore, $\tau^Q(\nu)$ is an optimal stopping time for maximizing $E_Q \left[Y_{\zeta} + \int_{\nu}^{\zeta} f(s, \theta_s^Q) ds \middle| \mathcal{F}_{\nu} \right]$ over $\zeta \in \mathcal{S}_{\nu,T}$.

For any $\nu \in \mathcal{S}_{0,T}$ and $k \in \mathbb{N}$, we introduce the collection of probability measures

$$\mathcal{Q}_{\nu}^k \triangleq \left\{ Q \in \mathcal{P}_{\nu} : |\theta_t^Q(\omega)| \vee f(t, \omega, \theta_t^Q(\omega)) \leq k, \quad dt \otimes dP\text{-a.e. on } \llbracket \nu, T \rrbracket \right\}.$$

Remark 3.2. It is clear that $\mathcal{Q}_{\nu}^k \subset \mathcal{Q}_{\nu}$; and from (f3) one can deduce that for any $\nu \leq \gamma$ we have

$$\mathcal{Q}_{\gamma} \subset \mathcal{Q}_{\nu} \quad \text{and} \quad \mathcal{Q}_{\gamma}^k \subset \mathcal{Q}_{\nu}^k, \quad \forall k \in \mathbb{N}.$$

Given a $Q \in \mathcal{Q}_{\nu}$ for some $\nu \in \mathcal{S}_{0,T}$, we *truncate* it in the following way: The predictability of process θ^Q and Proposition 2.1 imply that $\{f(t, \theta_t^Q)\}_{t \in [0,T]}$ is also a predictable process. Therefore, for any given $k \in \mathbb{N}$, the set

$$A_{\nu,k}^Q \triangleq \left\{ (t, \omega) \in \llbracket \nu, T \rrbracket : |\theta_t^Q(\omega)| \vee f(t, \omega, \theta_t^Q(\omega)) \leq k \right\} \in \mathcal{P} \quad (3.5)$$

is predictable. Then the predictable process $\theta^{Q^{\nu,k}} \triangleq \mathbf{1}_{A_{\nu,k}^Q} \theta^Q$ gives rise to a probability measure $Q^{\nu,k} \in \mathcal{Q}_{\nu}^k$ via the recipe $dQ^{\nu,k} \triangleq \mathcal{E}(\theta^{Q^{\nu,k}} \bullet B)_T dP$. Let us define the stopping times

$$\sigma_m^Q \triangleq \inf \left\{ t \in [0, T] : \int_0^t |\theta_s^Q|^2 ds > m \right\} \wedge T, \quad m \in \mathbb{N}.$$

There exists a null set N such that, for any $\omega \in \Omega \setminus N$, we have $\sigma_m^Q(\omega) = T$ for some $m = m(\omega) \in \mathbb{N}$. Since $E \int_0^{\sigma_m^Q} |\theta_t^Q|^2 dt \leq m$ holds for each $m \in \mathbb{N}$, we have $|\theta_t^Q(\omega)| < \infty$, $dt \otimes dP$ -a.e. on $\llbracket 0, \sigma_m^Q \rrbracket$.

As $\left(\bigcup_{m \in \mathbb{N}} \llbracket 0, \sigma_m^Q \rrbracket \right) \cup ([0, T] \times N) = [0, T] \times \Omega$, it follows that $|\theta_t^Q(\omega)| < \infty$ holds $dt \otimes dP$ -a.e. on $[0, T] \times \Omega$.

On the other hand, since $Q \in \mathcal{Q}_{\nu}$ we have $E_Q \int_{\nu}^T f(s, \theta_s^Q) ds < \infty$, which implies $\mathbf{1}_{\llbracket \nu, T \rrbracket}(t, \omega) f(t, \omega, \theta_t^Q(\omega)) < \infty$ holds $dt \otimes dQ$ -a.s., or equivalently $dt \otimes dP$ -a.e. Therefore, we see that

$$\lim_{k \rightarrow \infty} \uparrow \mathbf{1}_{A_{\nu,k}^Q} = \mathbf{1}_{\llbracket \nu, T \rrbracket}, \quad dt \otimes dP\text{-a.e.} \quad (3.6)$$

For any $\nu \in \mathcal{S}_{0,T}$, the upper value $\overline{V}(\nu)$ can be approximated from above in two steps, presented in the next two lemmas.

Lemma 3.1. Let $\nu \in \mathcal{S}_{0,T}$. (1) For any $\gamma \in \mathcal{S}_{\nu,T}$ we have

$$\operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\nu}} E_Q \left[Y_{\gamma} + \int_{\nu}^{\gamma} f(s, \theta_s^Q) ds \middle| \mathcal{F}_{\nu} \right] = \lim_{k \rightarrow \infty} \downarrow \operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\nu}^k} E_Q \left[Y_{\gamma} + \int_{\nu}^{\gamma} f(s, \theta_s^Q) ds \middle| \mathcal{F}_{\nu} \right], \quad P\text{-a.s.} \quad (3.7)$$

(2) It holds P -a.s. that

$$\overline{V}(\nu) = \operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\nu}} R^Q(\nu) = \lim_{k \rightarrow \infty} \downarrow \operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\nu}^k} R^Q(\nu). \quad (3.8)$$

Lemma 3.2. Let $k \in \mathbb{N}$ and $\nu \in \mathcal{S}_{0,T}$.

(1) For any $\gamma \in \mathcal{S}_{\nu,T}$ there exists a sequence $\{Q_n^{\gamma,k}\}_{n \in \mathbb{N}} \subset \mathcal{Q}_{\nu}^k$ such that

$$\operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\nu}^k} E_Q \left[Y_{\gamma} + \int_{\nu}^{\gamma} f(s, \theta_s^Q) ds \middle| \mathcal{F}_{\nu} \right] = \lim_{n \rightarrow \infty} \downarrow E_{Q_n^{\gamma,k}} \left[Y_{\gamma} + \int_{\nu}^{\gamma} f(s, \theta_s^{Q_n^{\gamma,k}}) ds \middle| \mathcal{F}_{\nu} \right], \quad P\text{-a.s.} \quad (3.9)$$

(2) There exists a sequence $\{Q_n^{(k)}\}_{n \in \mathbb{N}} \subset \mathcal{Q}_{\nu}^k$ such that

$$\operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\nu}^k} R^Q(\nu) = \lim_{n \rightarrow \infty} \downarrow R^{Q_n^{(k)}}(\nu), \quad P\text{-a.s.} \quad (3.10)$$

Let us fix $\nu \in \mathcal{S}_{0,T}$. For any $k \in \mathbb{N}$, the infimum of the family $\{\tau^Q(\nu)\}_{Q \in \mathcal{Q}_\nu^k}$ of optimal stopping times can be approached by a decreasing sequence in this family. As a result, the infimum is also a stopping time.

Lemma 3.3. *Let $\nu \in \mathcal{S}_{0,T}$ and $k \in \mathbb{N}$. There exists a sequence $\{Q_n^{(k)}\}_{n \in \mathbb{N}} \subseteq \mathcal{Q}_\nu^k$ such that*

$$\tau_k(\nu) \triangleq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} \tau^Q(\nu) = \lim_{n \rightarrow \infty} \downarrow \tau^{Q_n^{(k)}}(\nu), \quad P\text{-a.s.}$$

in the notation of Proposition 3.1, thus $\tau_k(\nu) \in \mathcal{S}_{\nu,T}$.

Since $\{\mathcal{Q}_\nu^k\}_{k \in \mathbb{N}}$ is an increasing sequence, $\{\tau_k(\nu)\}_{k \in \mathbb{N}}$ is in turn a decreasing sequence. Hence

$$\tau(\nu) \triangleq \lim_{k \rightarrow \infty} \downarrow \tau_k(\nu) \quad (3.11)$$

defines a stopping time in $\mathcal{S}_{\nu,T}$. The family of stopping times $\{\tau(\nu)\}_{\nu \in \mathcal{S}_{0,T}}$ will play a crucial role in this section.

The next lemma is concerned with the *pasting* of two probability measures.

Lemma 3.4. *Given $\nu \in \mathcal{S}_{0,T}$, let $\tilde{Q} \in \mathcal{Q}_\nu^k$ for some $k \in \mathbb{N}$. For any $Q \in \mathcal{Q}_\nu$ and $\gamma \in \mathcal{S}_{\nu,T}$, the predictable process*

$$\theta_t^{Q'} \triangleq \mathbf{1}_{\{t \leq \gamma\}} \theta_t^Q + \mathbf{1}_{\{t > \gamma\}} \theta_t^{\tilde{Q}}, \quad t \in [0, T] \quad (3.12)$$

induces a probability measure $Q' \in \mathcal{Q}_\nu$ by $dQ' \triangleq \mathcal{E}(\theta^{Q'} \bullet B)_T dP$. If Q belongs to $\mathcal{Q}_\nu^k$, so does Q' . Moreover, for any $\sigma \in \mathcal{S}_{\gamma,T}$, we have

$$R_\sigma^{Q',0} = R^{Q'}(\sigma) = R^{\tilde{Q}}(\sigma) = R_\sigma^{\tilde{Q},0}, \quad P\text{-a.s.} \quad (3.13)$$

Remark 3.3. *The probability measure Q' in Lemma 3.4 is called the pasting of Q and $\tilde{Q}$; see e.g. Section 6.7 of Föllmer and Schied [2004]. In general, $\mathcal{Q}_\nu$ is not closed under such “pasting”.*

The proofs of the following results use schemes similar to the ones in Karatzas and Zamfirescu [2008]. The main technical difficulty in our case is mentioned in Remark 3.3. Moreover, in order to use the results of Karatzas and Zamfirescu [2008] directly, we would have to assume that f and the θ^Q 's are all bounded. We overcome these difficulties by using approximation arguments that rely on *truncation* and *localization* techniques.

First, we shall show that at any $\nu \in \mathcal{S}_{0,T}$ we have $\underline{V}(\nu) = \overline{V}(\nu)$, P -a.s.

Theorem 3.1. Existence of Value: *For any $\nu \in \mathcal{S}_{0,T}$, we have*

$$\underline{V}(\nu) = \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_{\tau(\nu)} + \int_\nu^{\tau(\nu)} f(s, \theta_s^Q) ds \mid \mathcal{F}_\nu \right] = \overline{V}(\nu) \geq Y_\nu, \quad P\text{-a.s.} \quad (3.14)$$

Therefore, the stopping time $\tau(\nu)$ of (3.11) is optimal for the robust optimal stopping problem (3.1) (i.e., attains the essential infimum there).

We shall denote the common value in (3.14) by $V(\nu)$ ($= \underline{V}(\nu) = \overline{V}(\nu)$).

Proposition 3.2. *For any $\nu \in \mathcal{S}_{0,T}$, we have $V(\tau(\nu)) = Y_{\tau(\nu)}$, P -a.s.*

Note that $\tau(\nu)$ may not be the first time after ν when the value process coincides with the reward process. Actually, since the value process $\{V(t)\}_{t \in [0,T]}$ is not necessarily right-continuous, the random time $\inf\{t \in [\nu, T] : V(t) = Y_t\}$ may not even be a stopping time. We address this issue in the next three results.

Proposition 3.3. *Given $\nu \in \mathcal{S}_{0,T}$, $Q \in \mathcal{Q}_\nu$, and $\gamma \in \mathcal{S}_{\nu,\tau(\nu)}$, we have*

$$E_Q \left[V(\gamma) + \int_\nu^\gamma f(s, \theta_s^Q) ds \mid \mathcal{F}_\nu \right] \geq V(\nu), \quad P\text{-a.s.} \quad (3.15)$$

Lemma 3.5. For any $\nu, \gamma, \sigma \in \mathcal{S}_{0,T}$, we have the P -a.s. equalities

$$\mathbf{1}_{\{\nu=\gamma\}} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \mid \mathcal{F}_\nu \right] = \mathbf{1}_{\{\nu=\gamma\}} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\gamma} E_Q \left[Y_{\sigma \vee \gamma} + \int_\gamma^{\sigma \vee \gamma} f(s, \theta_s^Q) ds \mid \mathcal{F}_\gamma \right] \quad (3.16)$$

and

$$\mathbf{1}_{\{\nu=\gamma\}} V(\nu) = \mathbf{1}_{\{\nu=\gamma\}} V(\gamma). \quad (3.17)$$

Next, we show that for any given $\nu \in \mathcal{S}_{0,T}$, the process $\{\mathbf{1}_{\{t \geq \nu\}} V(\tau(\nu) \wedge t)\}_{t \in [0,T]}$ admits an RCLL modification $V^{0,\nu}$. As a consequence, the first time after ν when the process $V^{0,\nu}$ coincides with the process Y , is an optimal stopping time for the robust optimal stopping problem (3.1).

Theorem 3.2. Regularity of the Value: Let us fix a stopping time $\nu \in \mathcal{S}_{0,T}$.

(1) The process $\{\mathbf{1}_{\{t \geq \nu\}} V(\tau(\nu) \wedge t)\}_{t \in [0,T]}$ admits an RCLL modification $V^{0,\nu}$ such that, for any $\gamma \in \mathcal{S}_{0,T}$:

$$V_\gamma^{0,\nu} = \mathbf{1}_{\{\gamma \geq \nu\}} V(\tau(\nu) \wedge \gamma), \quad P\text{-a.s.} \quad (3.18)$$

(2) Consequently,

$$\tau_V(\nu) \triangleq \inf \left\{ t \in [\nu, T] : V_t^{0,\nu} = Y_t \right\} \quad (3.19)$$

is a stopping time which, in fact, attains the essential infimum in (3.1).

We should point out that, in order to determine the optimal stopping time in (1.1), knowledge of the function f in the representation (2.1) is not necessary. Indeed, let the ρ -Snell envelope be the RCLL modification of $\operatorname{ess\,sup}_{\gamma \in \mathcal{S}_{\nu,T}} (-\rho_{\nu,\gamma}(Y_\gamma))$, $\nu \in \mathcal{S}_{0,T}$. From our results above, the first time after ν that the ρ -Snell envelope touches the reward process Y is an optimal stopping time; this is consistent with the classical theory of optimal stopping.

4 The Saddle Point Problem

In this section we will construct a saddle point of the stochastic game in (1.2). As in the previous section, we shall assume here that $f : [0, T] \times \Omega \times \mathbb{R}^d \rightarrow [0, \infty]$ is a $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d) / \mathcal{B}([0, \infty])$ -measurable function which satisfies (f3). For any given $Q \in \mathcal{Q}_0$ and $\nu \in \mathcal{S}_{0,T}$, let us denote

$$Y_\nu^Q \triangleq Y_\nu + \int_0^\nu f(s, \theta_s^Q) ds \quad \text{and} \quad V^Q(\nu) \triangleq V(\nu) + \int_0^\nu f(s, \theta_s^Q) ds.$$

Definition 4.1. A pair $(Q^*, \sigma_*) \in \mathcal{Q}_0 \times \mathcal{S}_{0,T}$ is called a saddle point, if for every $Q \in \mathcal{Q}_0$ and $\nu \in \mathcal{S}_{0,T}$ we have

$$E_{Q^*}(Y_\nu^{Q^*}) \leq E_{Q^*}(Y_{\sigma_*}^{Q^*}) \leq E_Q(Y_{\sigma_*}^Q). \quad (4.1)$$

Theorem 4.1. Necessary Conditions for a Saddle Point: A pair $(Q^*, \sigma_*) \in \mathcal{Q}_0 \times \mathcal{S}_{0,T}$ is a saddle point, if the following conditions are satisfied:

- (i) $Y_{\sigma_*} = R^{Q^*}(\sigma_*)$, P -a.s.;
- (ii) for any $Q \in \mathcal{Q}_0$, we have $V(0) \leq E_Q[V^Q(\sigma_*)]$;
- (iii) for any $\nu \in \mathcal{S}_{0,\sigma_*}$, we have $V^{Q^*}(\nu) = E_{Q^*}[V^{Q^*}(\sigma_*) \mid \mathcal{F}_\nu]$, P -a.s.

To construct a saddle point, we need the following two notions.

Definition 4.2. We call $Z \in \widehat{\mathbb{H}}_{\mathbf{F}}^2([0, T]; \mathbb{R}^d)$ a BMO (short for Bounded Mean Oscillation) process if

$$\|Z\|_{BMO} \triangleq \sup_{\tau \in \mathcal{M}_{0,T}} \left\| E \left[\int_\tau^T |Z_s|^2 ds \mid \mathcal{F}_\tau \right]^{1/2} \right\|_\infty < \infty.$$

When Z is a BMO process, $Z \bullet B$ is a BMO martingale; see e.g. Kazamaki [1994].

Definition 4.3. BSDE with Reflection: Let $h : [0, T] \times \Omega \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$ be a $\widehat{\mathcal{P}} \times \mathcal{B}(\mathbb{R}) \times \mathcal{B}(\mathbb{R}^d)/\mathcal{B}(\mathbb{R})$ -measurable function. Given $S \in \mathbb{C}_{\mathbf{F}}^0[0, T]$ and $\xi \in \mathbb{L}^0(\mathcal{F}_T)$ with $\xi \geq S_T$, P -a.s., a triple $(\Gamma, \mathcal{Z}, K) \in \mathbb{C}_{\mathbf{F}}^0[0, T] \times \widehat{\mathbb{H}}_{\mathbf{F}}^2([0, T]; \mathbb{R}^d) \times \mathbb{K}_{\mathbf{F}}[0, T]$ is called a solution to the reflected backward stochastic differential equation with terminal condition ξ , generator h , and obstacle S (RBSDE (ξ, h, S) for short), if P -a.s., we have the comparison

$$S_t \leq \Gamma_t = \xi + \int_t^T h(s, \Gamma_s, \mathcal{Z}_s) ds + K_T - K_t - \int_t^T \mathcal{Z}_s dB_s, \quad t \in [0, T],$$

and the so-called flat-off condition

$$\int_0^T \mathbf{1}_{\{\Gamma_s > S_s\}} dK_s = 0, \quad P\text{-a.s.}$$

In the rest of this section we shall assume that the reward process $Y \in \mathbb{L}_{\mathbf{F}}^\infty[0, T]$ is continuous and that the function $f : [0, T] \times \Omega \times \mathbb{R}^d \rightarrow [0, \infty]$ satisfies the following additional conditions:

(H1) For every $(t, \omega) \in [0, T] \times \Omega$, the mapping $z \mapsto f(t, \omega, z)$ is continuous.

(H2) It holds $dt \otimes dP$ -a.e. that

$$f(t, \omega, z) \geq \varepsilon |z - \Upsilon_t(\omega)|^2 - \ell, \quad \forall z \in \mathbb{R}^d.$$

Here $\varepsilon > 0$ is a real constant, Υ is an $\mathbb{R}^d$ -valued process with $\|\Upsilon\|_\infty \triangleq \operatorname{ess\,sup}_{(t, \omega) \in [0, T] \times \Omega} |\Upsilon_t(\omega)| < \infty$, and $\ell \geq \varepsilon \|\Upsilon\|_\infty^2$.

(H3) For any $(t, \omega, u) \in [0, T] \times \Omega \times \mathbb{R}^d$, the mapping $z \mapsto f(t, \omega, z) + \langle u, z \rangle$ attains its infimum over $\mathbb{R}^d$ at some $z^* = z^*(t, \omega, u) \in \mathbb{R}^d$, namely,

$$\tilde{f}(t, \omega, u) \triangleq \inf_{z \in \mathbb{R}^d} (f(t, \omega, z) + \langle u, z \rangle) = f(t, \omega, z^*(t, \omega, u)) + \langle u, z^*(t, \omega, u) \rangle. \quad (4.2)$$

Without loss of generality, we can assume that the mapping $z^* : [0, T] \times \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d)/\mathcal{B}(\mathbb{R}^d)$ -measurable thanks to the Measurable Selection Theorem (see e.g. Lemma 1 of Beneš [1970] or Lemma 16.34 of Elliott [1982]). We further assume that there exist a non-negative BMO process ψ and a $M > 0$ such that for $dt \otimes dP$ -a.e. $(t, \omega) \in [0, T] \times \Omega$

$$|z^*(t, \omega, u)| \leq \psi_t(\omega) + M|u|, \quad \forall u \in \mathbb{R}^d.$$

Example 4.1. Let $\lambda \geq 0$ and let $\Lambda, \Upsilon \in \mathbb{H}_{\mathbf{F}}^\infty([0, T]; \mathbb{R}^d)$ with $\Lambda_t(\omega) \geq \varepsilon > 0$, $dt \otimes dP$ -a.e. Define

$$f(t, \omega, z) \triangleq \Lambda_t(\omega) \left(|z - \Upsilon_t(\omega)|^{2+\lambda} - |\Upsilon_t(\omega)|^{2+\lambda} \right), \quad \forall (t, \omega, z) \in [0, T] \times \Omega \times \mathbb{R}^d.$$

Clearly, $f^+ = f \vee 0$ is a $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d)/\mathcal{B}([0, \infty])$ -measurable function that satisfies (f3) and **(H1)**. It turns out that f^+ satisfies **(H2)**, since $dt \otimes dP$ -a.e. we have that

$$\begin{aligned} f^+(t, \omega, z) &\geq f(t, \omega, z) \geq \Lambda_t(\omega) \left(|z - \Upsilon_t(\omega)|^2 - 1 \right) - \Lambda_t(\omega) |\Upsilon_t(\omega)|^{2+\lambda} \\ &\geq \varepsilon |z - \Upsilon_t(\omega)|^2 - \|\Lambda\|_\infty (1 + \|\Upsilon\|_\infty^{2+\lambda}), \quad \forall z \in \mathbb{R}^d. \end{aligned}$$

For any $(t, \omega, u) \in [0, T] \times \Omega \times \mathbb{R}^d$ the gradient

$$\nabla_z (f(t, \omega, z) + \langle u, z \rangle) = (2 + \lambda) \Lambda_t(\omega) |z - \Upsilon_t(\omega)|^\lambda (z - \Upsilon_t(\omega)) + u, \quad \forall z \in \mathbb{R}^d,$$

is null only at $\hat{z}(t, \omega, u) = -[(2 + \lambda) \Lambda_t(\omega)]^{-\frac{1}{1+\lambda}} |u|^{-\frac{\lambda}{1+\lambda}} u + \Upsilon_t(\omega)$, where the mapping $z \mapsto f(t, \omega, z) + \langle u, z \rangle$ attains its infimum over $\mathbb{R}^d$. When $|u| \geq r_t(\omega) \triangleq (2 + \lambda) \Lambda_t(\omega) |\Upsilon_t(\omega)|^{1+\lambda}$, $\hat{z}(t, \omega, u) \in A \triangleq \{z \in \mathbb{R}^d : |z - \Upsilon_t(\omega)| \geq |\Upsilon_t(\omega)|\}$. It follows that

$$\begin{aligned} \inf_{z \in \mathbb{R}^d} (f^+(t, \omega, z) + \langle u, z \rangle) &\leq f^+(t, \omega, \hat{z}(t, \omega, u)) + \langle u, \hat{z}(t, \omega, u) \rangle = f(t, \omega, \hat{z}(t, \omega, u)) + \langle u, \hat{z}(t, \omega, u) \rangle \\ &= \inf_{z \in \mathbb{R}^d} (f(t, \omega, z) + \langle u, z \rangle) \leq \inf_{z \in \mathbb{R}^d} (f^+(t, \omega, z) + \langle u, z \rangle). \end{aligned} \quad (4.3)$$

On the other hand, when $|u| < r_t(\omega)$ or equivalently $\hat{z}(t, \omega, u) \notin A$, the gradient $\nabla_z(f(t, \omega, z) + \langle u, z \rangle) \neq 0$ for any $z \in A$, which implies that the mapping $z \rightarrow f(t, \omega, z) + \langle u, z \rangle$ can not attain its infimum over A at an interior point of it. Thus

$$\inf_{z \in A} (f(t, \omega, z) + \langle u, z \rangle) = \inf_{z \in \partial A} (f(t, \omega, z) + \langle u, z \rangle) = \inf_{z \in \partial A} \langle u, z \rangle.$$

Then it follows that

$$\inf_{z \in \mathbb{R}^d} (f^+(t, \omega, z) + \langle u, z \rangle) = \inf_{z \in A^c} \langle u, z \rangle \wedge \inf_{z \in A} (f(t, \omega, z) + \langle u, z \rangle) = \inf_{z \in \overline{A^c}} \langle u, z \rangle.$$

The latter infimum is attained uniquely at some $\tilde{z}(t, \omega, u) \in \overline{A^c}$, which together with (4.3) implies that

$$z^*(t, \omega, u) = \mathbf{1}_{\{|u| \geq r_t(\omega)\}} \hat{z}(t, \omega, u) + \mathbf{1}_{\{|u| < r_t(\omega)\}} \tilde{z}(t, \omega, u).$$

Therefore, f^+ satisfies (H3), since for $dt \otimes dP$ -a.e. $(t, \omega) \in [0, T] \times \Omega$ we have

$$\begin{aligned} |z^*(t, \omega, u)| &\leq |\hat{z}(t, \omega, u)| + |\tilde{z}(t, \omega, u)| \leq ((2 + \lambda)\varepsilon)^{-\frac{1}{1+\lambda}} |u|^{\frac{1}{1+\lambda}} + 3\|\Upsilon\|_\infty \\ &\leq ((2 + \lambda)\varepsilon)^{-\frac{1}{1+\lambda}} |u| + ((2 + \lambda)\varepsilon)^{-\frac{1}{1+\lambda}} + 3\|\Upsilon\|_\infty, \quad \forall u \in \mathbb{R}^d. \end{aligned}$$

Remark 4.1. The “entropic” risk measure with *risk tolerance coefficient* $r > 0$, namely

$$\rho_{\nu, \gamma}^r(\xi) \triangleq r \log \left\{ E \left[e^{-\frac{1}{r}\xi} \mid \mathcal{F}_\gamma \right] \right\}, \quad \xi \in \mathbb{L}^\infty(\mathcal{F}_\gamma),$$

is a typical example of a dynamic convex risk measures satisfying (A1)-(A4). The corresponding f in (2.1) is $f(z) = \frac{r}{2}|z|^2$, $z \in \mathbb{R}^d$.

Example 4.2. Let b^1, b^2 be two real-valued processes such that $-\varpi \leq b_t^1(\omega) \leq 0 \leq b_t^2(\omega) \leq \varpi$, $dt \otimes dP$ -a.e. for some $\varpi > 0$. Let $\varphi : [0, T] \times \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ be a $\mathcal{P} \otimes \mathcal{B}(\mathbb{R})/\mathcal{B}(\mathbb{R})$ -measurable function that satisfies the following two assumptions:

(i) For any $(t, \omega) \in [0, T] \times \Omega$, $\varphi(t, \omega, \cdot)$ is a bijective locally-integrable function or a continuous surjective locally-integrable function on $\mathbb{R}$.

(ii) For some $\varepsilon_1, \varepsilon_2 > 0$, it holds $dt \otimes dP$ -a.e. that

$$\varphi(t, \omega, x) \begin{cases} \geq (2\varepsilon_1 x + b_t^1(\omega)) \vee 0, & \text{if } x > 0, \\ \leq (2\varepsilon_2 x + b_t^2(\omega)) \wedge 0, & \text{if } x < 0. \end{cases}$$

Then $f(t, \omega, z) \triangleq \int_0^z \varphi(t, \omega, x) dx$, $z \in \mathbb{R}$ defines a $\mathcal{P} \otimes \mathcal{B}(\mathbb{R})/\mathcal{B}([0, \infty])$ -measurable non-negative function that satisfies (f3) and (H1). Let $\varepsilon = \varepsilon_1 \wedge \varepsilon_2$. For $dt \otimes dP$ -a.e. $(t, \omega) \in [0, T] \times \Omega$, if $z > 0$, then

$$f(t, \omega, z) \geq \int_0^z (2\varepsilon_1 x + b_t^1(\omega)) dx = \varepsilon_1 z^2 + b_t^1(\omega)z \geq \varepsilon z^2 - \varpi z = \varepsilon \left(z - \frac{\varpi}{2\varepsilon} \right)^2 - \frac{\varpi^2}{4\varepsilon};$$

on the other hand, if $z < 0$, then

$$\begin{aligned} f(t, \omega, z) &= - \int_z^0 \varphi(t, \omega, x) dx \geq - \int_z^0 (2\varepsilon_2 x + b_t^2(\omega)) dx = \varepsilon_2 z^2 + b_t^2(\omega)z \geq \varepsilon z^2 + \varpi z \\ &= \frac{1}{2}\varepsilon \left(z - \frac{\varpi}{2\varepsilon} \right)^2 + \frac{1}{2}\varepsilon \left(z + \frac{3\varpi}{2\varepsilon} \right)^2 - \frac{5\varpi^2}{4\varepsilon}. \end{aligned}$$

Thus it holds $dt \otimes dP$ -a.e. that $f(t, \omega, z) \geq \frac{1}{2}\varepsilon \left(z - \frac{\varpi}{2\varepsilon} \right)^2 - \frac{5\varpi^2}{4\varepsilon}$, i.e., (H2) is satisfied.

For any $(t, \omega, u) \in [0, T] \times \Omega \times \mathbb{R}$, since $\frac{d}{dz}(f(t, \omega, z) + uz) = \varphi(t, \omega, z) + u$, the mapping $z \mapsto f(t, \omega, z) + uz$ attains its infimum over $\mathbb{R}$ at each $z \in \{z \in \mathbb{R} : \varphi(t, \omega, z) = x\}$. Thus $\varphi_-^{-1}(t, \omega, x) \leq z^*(t, \omega, u) \leq \varphi_+^{-1}(t, \omega, x)$, where

$$\varphi_-^{-1}(t, \omega, x) \triangleq \inf\{z \in \mathbb{R} : \varphi(t, \omega, z) = x\} \quad \text{and} \quad \varphi_+^{-1}(t, \omega, x) \triangleq \sup\{z \in \mathbb{R} : \varphi(t, \omega, z) = x\}.$$

It is clear that $\varphi(t, \omega, \varphi_-^{-1}(t, \omega, x)) = x$ and $\varphi(t, \omega, \varphi_+^{-1}(t, \omega, x)) = x$. For $dt \otimes dP$ -a.e. $(t, \omega) \in [0, T] \times \Omega$ and $u \in \mathbb{R}$, if $\varphi_-^{-1}(t, \omega, x) > 0$, then

$$-u = \varphi(t, \omega, \varphi_-^{-1}(t, \omega, -u)) \geq 2\varepsilon_1 \varphi_-^{-1}(t, \omega, x) + b_t^1(\omega),$$

which implies that $0 < \varphi_-^{-1}(t, \omega, x) \leq \frac{1}{2\varepsilon}(|u| + \varpi)$. On the other hand, if $\varphi_-^{-1}(t, \omega, x) < 0$, one can deduce that $-\frac{1}{2\varepsilon}(|u| + \varpi) \leq \varphi_-^{-1}(t, \omega, x) < 0$ by a similar argument. Hence $|\varphi_-^{-1}(t, \omega, x)| \leq \frac{1}{2\varepsilon}(|u| + \varpi)$. Similarly, this inequality also holds for $\varphi_+^{-1}(t, \omega, x)$, thus for $z^*(t, \omega, u)$. As a result, (H3) is also satisfied. $\square$

One can easily deduce from **(H2)** and (f3) that $dt \otimes dP$ -a.e.

$$-\frac{1+\varepsilon}{4\varepsilon}|u|^2 - \|\Upsilon\|_\infty^2 - \ell \leq \tilde{f}(t, \omega, u) \leq 0, \quad \forall u \in \mathbb{R}^d,$$

which shows that $\tilde{f}$ has quadratic growth in u . Thanks to Theorems 1 and 3 of Kobylanski et al. [2002], the RBSDE $(Y_T, \tilde{f}, Y)$ admits a solution $(\tilde{\Gamma}, \tilde{Z}, \tilde{K}) \in \mathbb{C}_F^\infty[0, T] \times \mathbb{H}_F^2([0, T]; \mathbb{R}^d) \times \mathbb{K}_F[0, T]$.

In fact, $\tilde{Z}$ is a BMO process. To see this, we set $\kappa \triangleq \frac{1+\varepsilon}{4\varepsilon} \vee (\|\Upsilon\|^2 + \ell)$. For any $\nu \in \mathcal{S}_{0,T}$, applying Itô's formula to $e^{-4\kappa\tilde{\Gamma}_t}$ we get

$$\begin{aligned} e^{-4\kappa\tilde{\Gamma}_\nu} + 8\kappa^2 \int_\nu^T e^{-4\kappa\tilde{\Gamma}_s} |\tilde{Z}_s|^2 ds &= e^{-4\kappa Y_T} - 4\kappa \int_\nu^T e^{-4\kappa\tilde{\Gamma}_s} \tilde{f}(s, \tilde{Z}_s) ds - 4\kappa \int_\nu^T e^{-4\kappa\tilde{\Gamma}_s} d\tilde{K}_s + 4\kappa \int_\nu^T e^{-4\kappa\tilde{\Gamma}_s} \tilde{Z}_s dB_s \\ &\leq e^{-4\kappa Y_T} + 4\kappa^2 \int_\nu^T e^{-4\kappa\tilde{\Gamma}_s} (1 + |\tilde{Z}_s|^2) ds + 4\kappa \int_\nu^T e^{-4\kappa\tilde{\Gamma}_s} \tilde{Z}_s dB_s. \end{aligned}$$

Taking conditional expectations in the above expression, we obtain

$$e^{-4\kappa\|\tilde{\Gamma}\|_\infty} E \left[\int_\nu^T |\tilde{Z}_s|^2 ds \mid \mathcal{F}_\nu \right] \leq E \left[\int_\nu^T e^{-4\kappa\tilde{\Gamma}_s} |\tilde{Z}_s|^2 ds \mid \mathcal{F}_\nu \right] \leq \frac{1}{4\kappa^2} E \left[e^{-4\kappa Y_T} \mid \mathcal{F}_\nu \right] + e^{4\kappa\|\tilde{\Gamma}\|_\infty} T.$$

which implies that $\|\tilde{Z}\|_{BMO} \leq e^{4\kappa\|\tilde{\Gamma}\|_\infty} \left(\frac{1}{4\kappa^2} + T \right)^{1/2}$.

Since the mapping $z^* : [0, T] \times \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d)/\mathcal{B}(\mathbb{R}^d)$ -measurable (see (H3)),

$$\theta_t^*(\omega) \triangleq z^*(t, \omega, \tilde{Z}_t(\omega)), \quad (t, \omega) \in [0, T] \times \Omega \quad (4.4)$$

is a predictable process. It follows from **(H3)** that for any $\nu \in [0, T]$

$$E \left[\int_\nu^T |\theta_s^*|^2 ds \mid \mathcal{F}_\nu \right] \leq 2E \left[\int_\nu^T \psi_s^2 ds \mid \mathcal{F}_\nu \right] + 2M^2 E \left[\int_\nu^T |\tilde{Z}_s|^2 ds \mid \mathcal{F}_\nu \right], \quad P\text{-a.s.},$$

which implies that θ^* is a BMO process.

Fix $\nu \in \mathcal{S}_{0,T}$. Since $\theta_t^{*,\nu} \triangleq \mathbf{1}_{\{t > \nu\}} \theta_t^*$, $t \in [0, T]$ is also a BMO process, we know from Theorem 2.3 of Kazamaki [1994] that the stochastic exponential $\{\mathcal{E}(\theta^{*,\nu} \bullet B)_t\}_{t \in [0, T]}$ is a uniformly integrable martingale. Therefore, $dQ^{*,\nu} \triangleq \mathcal{E}(\theta^{*,\nu} \bullet B)_T dP$ defines a probability measure $Q^{*,\nu} \in \mathcal{P}_\nu$. As

$$\tilde{f}(s, \tilde{Z}_s) = f(s, z^*(s, \tilde{Z}_s)) + \langle \tilde{Z}_s, z^*(s, \tilde{Z}_s) \rangle = f(s, \theta_s^*) + \langle \tilde{Z}_s, \theta_s^* \rangle, \quad dt \otimes dP\text{-a.e.}$$

by (4.2) and (4.4) and the Girsanov Theorem, we can deduce

$$\begin{aligned} \tilde{\Gamma}_{\nu \vee t} &= Y_T + \int_{\nu \vee t}^T \left[f(s, \theta_s^{*,\nu}) + \langle \tilde{Z}_s, \theta_s^{*,\nu} \rangle \right] ds + \tilde{K}_T - \tilde{K}_{\nu \vee t} - \int_{\nu \vee t}^T \tilde{Z}_s dB_s \\ &= Y_T + \int_{\nu \vee t}^T f(s, \theta_s^{*,\nu}) ds + \tilde{K}_T - \tilde{K}_{\nu \vee t} - \int_{\nu \vee t}^T \tilde{Z}_s dB_s^{Q^{*,\nu}}, \quad t \in [0, T], \end{aligned} \quad (4.5)$$

where $B^{Q^{*,\nu}}$ is a Brownian Motion under $Q^{*,\nu}$. Letting $t = 0$ and taking the expectation $E_{Q^{*,\nu}}$ yield that

$$E_{Q^{*,\nu}} \int_\nu^T f(s, \theta_s^{*,\nu}) ds \leq E_{Q^{*,\nu}} (\tilde{\Gamma}_\nu - Y_T) \leq 2\|\tilde{\Gamma}\|_\infty,$$

thus $Q^{*,\nu} \in \mathcal{Q}_\nu$. The lemma below shows that $\tilde{\Gamma}$ is indistinguishable from $R^{Q^{*,\nu}, 0}$ on the stochastic interval $[\nu, T]$.

Lemma 4.1. *Given $\nu \in \mathcal{S}_{0,T}$, it holds P -a.s. that*

$$\tilde{\Gamma}_t = R_t^{Q^{*,\nu},0}, \quad \forall t \in [\nu, T]. \quad (4.6)$$

Let $k \in \mathbb{N}$ and $Q \in \mathcal{Q}_\nu^k$. It is easy to see that the function $h_Q(s, \omega, z) \triangleq f(s, \omega, \theta_s^Q(\omega)) + \langle z, \theta_s^Q(\omega) \rangle$ is Lipschitz continuous in z : to wit, for $dt \otimes dP$ -a.e. $(t, \omega) \in [0, T] \times \Omega$

$$|h_Q(s, \omega, z) - h_Q(s, \omega, z')| = |\langle z - z', \theta_s^Q \rangle| \leq |\theta_s^Q| \cdot |z - z'| \leq k|z - z'|, \quad \forall z, z' \in \mathbb{R}^d.$$

Moreover, we have

$$E \int_0^T |h_Q(s, 0)|^2 ds = E \int_0^T |f(s, \theta_s^Q)|^2 ds = E \int_\nu^T |f(s, \theta_s^Q)|^2 ds \leq k^2 T.$$

Theorem 5.2 of El Karoui et al. [1997] assures now that there exists a unique solution $(\Gamma^Q, \mathcal{Z}^Q, K^Q) \in \mathbb{C}_{\mathbf{F}}^2[0, T] \times \mathbb{H}_{\mathbf{F}}^2([0, T]; \mathbb{R}^d) \times \mathbb{K}_{\mathbf{F}}[0, T]$ to the RBSDE (Y_T, h_Q, Y) . Fix $t \in [0, T]$. For any $\gamma \in \mathcal{S}_{t,T}$, Girsanov Theorem implies

$$\begin{aligned} \Gamma_t^Q &= Y_T + \int_t^T h_Q(s, \mathcal{Z}_s^Q) ds + K_T^Q - K_t^Q - \int_t^T \mathcal{Z}_s^Q dB_s \\ &= \Gamma_\gamma^Q + \int_t^\gamma f(s, \theta_s^Q) ds + K_\gamma^Q - K_t^Q - \int_t^\gamma \mathcal{Z}_s^Q dB_s^Q, \quad P\text{-a.s.}, \end{aligned}$$

where B^Q is a Brownian Motion under Q . By analogy with Lemma 4.1, it holds P -a.s. that

$$\Gamma_t^Q = R_t^{Q,0}, \quad \forall t \in [0, T]. \quad (4.7)$$

In particular, we see that $R^{Q,0}$ is, in fact, a continuous process.

Next, we recall a comparison theorem of RBSDEs; see Theorem 4.1 of El Karoui et al. [1997]. (We restate it in a more general form.)

Proposition 4.1. *Let $(\Gamma, \mathcal{Z}, K)$ (resp. $(\Gamma', \mathcal{Z}', K')$) be a solution of RBSDE (ξ, h, S) (resp. RBSDE (ξ', h', S')) in the sense of Definition 4.3. Additionally, assume that*

- (i) *either h or h' is Lipschitz in (y, z) ;*
- (ii) *it holds P -a.s. that $\xi \leq \xi'$ and $S_t \leq S'_t$ for any $t \in [0, T]$;*
- (iii) *it holds $dt \otimes dP$ -a.e. that $h(t, \omega, y, z) \leq h'(t, \omega, y, z)$ for any $(y, z) \in \mathbb{R} \times \mathbb{R}^d$.*

Then it holds P -a.s. that $\Gamma_t \leq \Gamma'_t$ for any $t \in [0, T]$.

Since it holds $dt \otimes dP$ -a.e. that

$$\tilde{f}(t, \omega, u) \triangleq \inf_{z \in \mathbb{R}^d} (f(t, \omega, z) + \langle u, z \rangle) \leq f(s, \omega, \theta_s^Q(\omega)) + \langle u, \theta_s^Q(\omega) \rangle = h_Q(s, \omega, u), \quad \forall u \in \mathbb{R}^d.$$

we see from Proposition 4.1 and (4.7) that we have P -a.s.

$$\tilde{\Gamma}_t \leq \Gamma_t^Q = R_t^{Q,0}, \quad \forall t \in [0, T]. \quad (4.8)$$

Letting $t = \nu$, taking the essential infimum of right-hand-side over $Q \in \mathcal{Q}_\nu^k$, and then letting $k \rightarrow \infty$, we can deduce from Lemma 4.1, (3.8), and (3.3) that

$$R_\nu^{Q^{*,\nu},0} = \tilde{\Gamma}_\nu \leq \lim_{k \rightarrow \infty} \downarrow \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} R_\nu^{Q,0} = \lim_{k \rightarrow \infty} \downarrow \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} R^Q(\nu) = \overline{V}(\nu) = V(\nu) \leq R^{Q^{*,\nu}}(\nu) = R_\nu^{Q^{*,\nu},0}, \quad P\text{-a.s.}$$

which implies that $V(\nu) = \tilde{\Gamma}_\nu$, P -a.s. Applying Lemma 4.1 and (3.3) once again yields that

$$V(\nu) = \tilde{\Gamma}_\nu = R_\nu^{Q^*,0} = R^{Q^*}(\nu), \quad P\text{-a.s.} \quad (4.9)$$

where $Q^* \triangleq Q^{*,0} \in \mathcal{Q}_0$. It is clear that $dQ^* = dQ^{*,0} = \mathcal{E}(\theta^{*,0} \bullet B)_T dP = \mathcal{E}(\theta^* \bullet B)_T dP$.

We are now ready to state the main result of this section.

Theorem 4.2. Existence of a Saddle Point: *The pair $(Q^*, \tau^{Q^*}(0))$ is a saddle point as in (4.1).*

5 Proofs

5.1 Proof of the Results in Sections 2 and 3

Proof of Proposition 2.1: Bion-Nadal [2009, Proposition 1] shows that

$$\rho_{\nu,\gamma}(\xi) = \operatorname{ess\,sup}_{Q \in \mathcal{Q}_{\nu,\gamma}} \left(E_Q [-\xi | \mathcal{F}_\nu] - \alpha_{\nu,\gamma}(Q) \right), \quad P\text{-a.s.} \quad (5.1)$$

Here we have set $\mathcal{Q}_{\nu,\gamma} \triangleq \{Q \in \mathcal{P}_\nu : E_Q[\alpha_{\nu,\gamma}(Q)] < \infty\}$, and the quantity

$$\alpha_{\nu,\gamma}(Q) \triangleq \operatorname{ess\,sup}_{\eta \in \mathbb{L}^\infty(\mathcal{F}_\gamma)} \left(E_Q [-\eta | \mathcal{F}_\nu] - \rho_{\nu,\gamma}(\eta) \right)$$

is known as the “minimal penalty” of $\rho_{\nu,\gamma}$. (The representation (5.1) was shown for $Q \ll P$ rather than $Q \sim P$ in Bion-Nadal [2009]. However, our assumption **(A4)** assures that (5.1) also holds. For a proof, see Föllmer and Penner [2006, Lemma 3.5] and Klöppel and Schweizer [2007, Theorem 3.1].)

Thanks to Delbaen et al. [2009, Theorem 5(i) and the proof of Proposition 9(v)], there exists a nonnegative function $f : [0, T] \times \Omega \times \mathbb{R}^d \rightarrow [0, \infty]$ satisfying (f1)-(f3), such that for each $Q \in \mathcal{Q}_{\nu,\gamma}$ we have

$$\alpha_{\nu,\gamma}(Q) = E_Q \left(\int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right), \quad P\text{-a.s.}$$

Hence we can rewrite $\mathcal{Q}_{\nu,\gamma} = \{Q \in \mathcal{P}_\nu : E_Q \int_\nu^\gamma f(s, \theta_s^Q) ds < \infty\}$, and (5.1) becomes

$$\rho_{\nu,\gamma}(\xi) = \operatorname{ess\,sup}_{Q \in \mathcal{Q}_{\nu,\gamma}} E_Q \left[-\xi - \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.} \quad (5.2)$$

Since $\mathcal{Q}_\nu \equiv \mathcal{Q}_{\nu,T} \subset \mathcal{Q}_{\nu,\gamma}$, it follows readily that

$$\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \geq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\nu,\gamma}} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.} \quad (5.3)$$

On the other hand, for any given $Q \in \mathcal{Q}_{\nu,\gamma}$, the predictable process $\theta_t^{\tilde{Q}} \triangleq \mathbf{1}_{\{t \leq \gamma\}} \theta_t^Q$, $t \in [0, T]$ induces a probability measure $\tilde{Q} \in \mathcal{P}_\nu$ via $d\tilde{Q} \triangleq \mathcal{E}(\theta^{\tilde{Q}} \bullet B)_T dP$. Since $f(t, \theta_t^{\tilde{Q}}) = \mathbf{1}_{\{t \leq \gamma\}} f(t, \theta_t^Q)$, $dt \otimes dP$ -a.e. from (f3), it follows

$$E_{\tilde{Q}} \int_\nu^T f(s, \theta_s^{\tilde{Q}}) ds = E_{\tilde{Q}} \int_\nu^\gamma f(s, \theta_s^{\tilde{Q}}) ds = E_Q \int_\nu^\gamma f(s, \theta_s^Q) ds < \infty,$$

thus $\tilde{Q} \in \mathcal{Q}_\nu$. Then we can deduce

$$\begin{aligned} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] &\leq E_{\tilde{Q}} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{\tilde{Q}}) ds \middle| \mathcal{F}_\nu \right] = E_{\tilde{Q}} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \\ &= E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.} \end{aligned}$$

Taking the essential infimum of the right-hand-side over $Q \in \mathcal{Q}_{\nu,\gamma}$ yields

$$\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \leq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\nu,\gamma}} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.};$$

this, together with (5.3) and (5.2), proves (2.1). $\square$

Proof of Lemma 3.1: (1) Since $\{\mathcal{Q}_\nu^k\}_{k \in \mathbb{N}}$ is an increasing sequence of sets contained in $\mathcal{Q}_\nu$, it follows that

$$\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \leq \lim_{k \rightarrow \infty} \downarrow \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.} \quad (5.4)$$

Now let us fix a probability measure $Q \in \mathcal{Q}_\nu$, and define the stopping times

$$\delta_m^Q \triangleq \inf \left\{ t \in [\nu, T] : \int_\nu^t [f(s, \theta_s^Q) + |\theta_s^Q|^2] ds > m \right\} \wedge T, \quad m \in \mathbb{N}.$$

It is easy to see that $\lim_{m \rightarrow \infty} \uparrow \delta_m^Q = T$, P -a.s. For any $(m, k) \in \mathbb{N}^2$, the predictable process $\theta_t^{Q^{m,k}} \triangleq \mathbf{1}_{\{t \leq \delta_m^Q\}} \mathbf{1}_{A_{\nu,k}^Q} \theta_t^Q$, $t \in [0, T]$ induces a probability measure $Q^{m,k} \in \mathcal{Q}_\nu^k$ by

$$dQ^{m,k} \triangleq \mathcal{E}(\theta^{Q^{m,k}} \bullet B)_T \cdot dP \quad (5.5)$$

(recall the notation of (3.5)). It follows from (f3) that

$$f(t, \theta_t^{Q^{m,k}}) = \mathbf{1}_{\{t \leq \delta_m^Q\}} \mathbf{1}_{A_{\nu,k}^Q} f(t, \theta_t^Q), \quad dt \otimes dP\text{-a.e.} \quad (5.6)$$

Then we can deduce from Bayes' Rule (see, e.g., Karatzas and Shreve [1991, Lemma 3.5.3]) that

$$\begin{aligned} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] &\leq E_{Q^{m,k}} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q^{m,k}}) ds \middle| \mathcal{F}_\nu \right] \\ &= E \left[Z_{\nu,T}^{Q^{m,k}} \left(Y_\gamma + \int_\nu^{\gamma \wedge \delta_m^Q} \mathbf{1}_{A_{\nu,k}^Q} f(s, \theta_s^Q) ds \right) \middle| \mathcal{F}_\nu \right] \leq E \left[Z_{\nu,T}^{Q^{m,k}} \left(Y_\gamma + \int_\nu^{\gamma \wedge \delta_m^Q} f(s, \theta_s^Q) ds \right) \middle| \mathcal{F}_\nu \right] \\ &= E \left[\left(Z_{\nu,T}^{Q^{m,k}} - Z_{\nu,\delta_m^Q}^Q \right) \left(Y_\gamma + \int_\nu^{\gamma \wedge \delta_m^Q} f(s, \theta_s^Q) ds \right) \middle| \mathcal{F}_\nu \right] + E \left[(Z_{\nu,\delta_m^Q}^Q - Z_{\nu,T}^Q) \cdot Y_\gamma \middle| \mathcal{F}_\nu \right] \\ &\quad + E \left[Z_{\nu,T}^Q Y_\gamma \middle| \mathcal{F}_\nu \right] + E \left[Z_{\nu,\delta_m^Q}^Q \int_\nu^{\gamma \wedge \delta_m^Q} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \\ &\leq (\|Y\|_\infty + m) \cdot E \left[\left| Z_{\nu,T}^{Q^{m,k}} - Z_{\nu,\delta_m^Q}^Q \right| \middle| \mathcal{F}_\nu \right] + \|Y\|_\infty \cdot E \left[\left| Z_{\nu,\delta_m^Q}^Q - Z_{\nu,T}^Q \right| \middle| \mathcal{F}_\nu \right] + E_Q \left[Y_\gamma \middle| \mathcal{F}_\nu \right] \\ &\quad + E_Q \left[\int_\nu^{\gamma \wedge \delta_m^Q} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \\ &\leq (\|Y\|_\infty + m) \cdot E \left[\left| Z_{\nu,T}^{Q^{m,k}} - Z_{\nu,\delta_m^Q}^Q \right| \middle| \mathcal{F}_\nu \right] + \|Y\|_\infty \cdot E \left[\left| Z_{\nu,\delta_m^Q}^Q - Z_{\nu,T}^Q \right| \middle| \mathcal{F}_\nu \right] \\ &\quad + E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.} \end{aligned} \quad (5.7)$$

From the equation (3.6) and the Dominated Convergence Theorem, we observe

$$\lim_{k \rightarrow \infty} E \left(\int_\nu^{\delta_m^Q} (\mathbf{1}_{A_{\nu,k}^Q} - 1) \theta_s^Q dB_s \right)^2 = \lim_{k \rightarrow \infty} E \int_\nu^{\delta_m^Q} (1 - \mathbf{1}_{A_{\nu,k}^Q}) |\theta_s^Q|^2 ds = 0, \quad P\text{-a.s.}$$

Thus we can find a subsequence of $\{A_{\nu,k}^Q\}_{k \in \mathbb{N}}$ (we still denote it by $\{A_{\nu,k}^Q\}_{k \in \mathbb{N}}$) such that

$$\lim_{k \rightarrow \infty} \int_\nu^{\delta_m^Q} \mathbf{1}_{A_{\nu,k}^Q} \theta_s^Q dB_s = \int_\nu^{\delta_m^Q} \theta_s^Q dB_s \quad \text{and} \quad \lim_{k \rightarrow \infty} \int_\nu^{\delta_m^Q} \mathbf{1}_{A_{\nu,k}^Q} |\theta_s^Q|^2 ds = \int_\nu^{\delta_m^Q} |\theta_s^Q|^2 ds, \quad P\text{-a.s.}$$

and consequently, P -a.s.:

$$\lim_{k \rightarrow \infty} Z_{\nu,T}^{Q^{m,k}} = \lim_{k \rightarrow \infty} \exp \left\{ \int_\nu^{\delta_m^Q} \mathbf{1}_{A_{\nu,k}^Q} \left(\theta_s^Q dB_s - \frac{1}{2} |\theta_s^Q|^2 ds \right) \right\} = \exp \left\{ \int_\nu^{\delta_m^Q} \left(\theta_s^Q dB_s - \frac{1}{2} |\theta_s^Q|^2 ds \right) \right\} = Z_{\nu,\delta_m^Q}^Q.$$

Since $E(Z_{\nu,T}^{Q^{m,k}} | \mathcal{F}_\nu) = E(Z_{\nu,\delta_m^Q}^Q | \mathcal{F}_\nu) = 1$, P -a.s. for any $k \in \mathbb{N}$, it follows from Scheffé's Lemma (see e.g. Williams [1991, Section 5.10]) that

$$\lim_{k \rightarrow \infty} E \left[\left| Z_{\nu,T}^{Q^{m,k}} - Z_{\nu,\delta_m^Q}^Q \right| \middle| \mathcal{F}_\nu \right] = 0, \quad P\text{-a.s.} \quad (5.8)$$

Hence, letting $k \rightarrow \infty$ in (5.7), we obtain

$$\begin{aligned} \lim_{k \rightarrow \infty} \downarrow \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \\ \leq E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] + \|Y\|_\infty \cdot E \left[|Z_{\nu, \delta_m^Q}^Q - Z_{\nu, T}^Q| \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.} \end{aligned} \quad (5.9)$$

It is easy to see that $\lim_{m \rightarrow \infty} \uparrow \delta_m^Q = T$, P -a.s. The right-continuity of the process Z^Q then implies that $\lim_{m \rightarrow \infty} Z_{\nu, \delta_m^Q}^Q = Z_{\nu, T}^Q$, P -a.s. Since $E \left[Z_{\nu, \delta_m^Q}^Q \middle| \mathcal{F}_\nu \right] = E \left[Z_{\nu, T}^Q \middle| \mathcal{F}_\nu \right] = 1$, P -a.s. for any $m \in \mathbb{N}$, using Scheffé's Lemma once again we obtain

$$\lim_{m \rightarrow \infty} E \left[|Z_{\nu, \delta_m^Q}^Q - Z_{\nu, T}^Q| \middle| \mathcal{F}_\nu \right] = 0, \quad P\text{-a.s.} \quad (5.10)$$

Therefore, letting $m \rightarrow \infty$ in (5.9) we obtain

$$\lim_{k \rightarrow \infty} \downarrow \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \leq E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.}$$

Taking the essential infimum of right-hand-side over $Q \in \mathcal{Q}_\nu$ gives

$$\lim_{k \rightarrow \infty} \downarrow \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \leq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.}$$

which, together with (5.4), proves (3.7).

(2) By analogy with (5.4), we have

$$\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} R^Q(\nu) \leq \lim_{k \rightarrow \infty} \downarrow \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} R^Q(\nu), \quad P\text{-a.s.} \quad (5.11)$$

Taking the essential supremum in (5.7) over $\gamma \in \mathcal{S}_{\nu, T}$ we get

$$\begin{aligned} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} R^Q(\nu) \leq R^{Q^{m, k}}(\nu) &\leq R^Q(\nu) + (\|Y\|_\infty + m) \cdot E \left[|Z_{\nu, T}^{Q^{m, k}} - Z_{\nu, \delta_m^Q}^Q| \middle| \mathcal{F}_\nu \right] \\ &\quad + \|Y\|_\infty \cdot E \left[|Z_{\nu, \delta_m^Q}^Q - Z_{\nu, T}^Q| \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.} \end{aligned} \quad (5.12)$$

In light of (5.8) and (5.10), letting $k \rightarrow \infty$ and subsequently letting $m \rightarrow \infty$ in (5.12), we obtain

$$\lim_{k \rightarrow \infty} \downarrow \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} R^Q(\nu) \leq R^Q(\nu), \quad P\text{-a.s.}$$

Taking the essential infimum of right-hand-side over $Q \in \mathcal{Q}_\nu$ yields $\lim_{k \rightarrow \infty} \downarrow \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} R^Q(\nu) \leq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} R^Q(\nu)$, P -a.s. which, together with (5.11), proves (3.8). $\square$

Proof of Lemma 3.2: (1) We first show that the family $\left\{ E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \right\}_{Q \in \mathcal{Q}_\nu^k}$ is directed downwards, i.e., for any $Q_1, Q_2 \in \mathcal{Q}_\nu^k$, there exists a $Q_3 \in \mathcal{Q}_\nu^k$ such that

$$E_{Q_3} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_3}) ds \middle| \mathcal{F}_\nu \right] \leq E_{Q_1} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_1}) ds \middle| \mathcal{F}_\nu \right] \wedge E_{Q_2} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_2}) ds \middle| \mathcal{F}_\nu \right] \quad P\text{-a.s.} \quad (5.13)$$

To see this, we let $Q_1, Q_2 \in \mathcal{Q}_\nu^k$ and let $A \in \mathcal{F}_\nu$. It is clear that

$$\theta_t^{Q_3} \triangleq \mathbf{1}_{\{t > \nu\}} \left(\mathbf{1}_A \theta_t^{Q_1} + \mathbf{1}_{A^c} \theta_t^{Q_2} \right), \quad t \in [0, T] \quad (5.14)$$

forms a predictable process, thus we can define a probability measure $Q_3 \in \mathcal{M}^e$ via $dQ_3 \triangleq \mathcal{E}(\theta^{Q_3} \bullet B)_T dP$. It follows from (f3) that

$$f(t, \theta_t^{Q_3}) = \mathbf{1}_{\{t > \nu\}} \left(\mathbf{1}_A f(t, \theta_t^{Q_1}) + \mathbf{1}_{A^c} f(t, \theta_t^{Q_2}) \right), \quad dt \otimes dP\text{-a.e.}, \quad (5.15)$$

which together with (5.14) implies that $\theta^{Q_3} = 0$ $dt \otimes dP$ -a.e. on $\llbracket 0, \nu \rrbracket$ and $|\theta_t^{Q_3}(\omega)| \vee f(t, \omega, \theta_t^{Q_3}(\omega)) = \mathbf{1}_A(\omega) |\theta_t^{Q_1}(\omega)| \vee f(t, \omega, \theta_t^{Q_1}(\omega)) + \mathbf{1}_{A^c}(\omega) |\theta_t^{Q_2}(\omega)| \vee f(t, \omega, \theta_t^{Q_2}(\omega)) \leq k$, $dt \otimes dP$ -a.e. on $\llbracket \nu, T \rrbracket$. Hence $Q_3 \in \mathcal{Q}_\nu^k$. For any $\gamma \in \mathcal{S}_{\nu, T}$, we have

$$\begin{aligned} Z_{\nu, \gamma}^{Q_3} &= \exp \left\{ \int_\nu^\gamma (\mathbf{1}_A \theta_s^{Q_1} + \mathbf{1}_{A^c} \theta_s^{Q_2}) dB_s - \frac{1}{2} \int_\nu^\gamma (\mathbf{1}_A |\theta_s^{Q_1}|^2 + \mathbf{1}_{A^c} |\theta_s^{Q_2}|^2) ds \right\} \\ &= \exp \left\{ \mathbf{1}_A \left(\int_\nu^\gamma \theta_s^{Q_1} dB_s - \frac{1}{2} \int_\nu^\gamma |\theta_s^{Q_1}|^2 ds \right) + \mathbf{1}_{A^c} \left(\int_\nu^\gamma \theta_s^{Q_2} dB_s - \frac{1}{2} \int_\nu^\gamma |\theta_s^{Q_2}|^2 ds \right) \right\} \\ &= \mathbf{1}_A \exp \left\{ \int_\nu^\gamma \theta_s^{Q_1} dB_s - \frac{1}{2} \int_\nu^\gamma |\theta_s^{Q_1}|^2 ds \right\} + \mathbf{1}_{A^c} \exp \left\{ \int_\nu^\gamma \theta_s^{Q_2} dB_s - \frac{1}{2} \int_\nu^\gamma |\theta_s^{Q_2}|^2 ds \right\} \\ &= \mathbf{1}_A Z_{\nu, \gamma}^{Q_1} + \mathbf{1}_{A^c} Z_{\nu, \gamma}^{Q_2}, \quad P\text{-a.s.} \end{aligned} \quad (5.16)$$

Then Bayes' Rule implies that

$$\begin{aligned} E_{Q_3} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_3}) ds \middle| \mathcal{F}_\nu \right] &= E \left[Z_{\nu, T}^{Q_3} \left(Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_3}) ds \right) \middle| \mathcal{F}_\nu \right] \\ &= E \left[\mathbf{1}_A Z_{\nu, T}^{Q_1} \left(Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_1}) ds \right) + \mathbf{1}_{A^c} Z_{\nu, T}^{Q_2} \left(Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_2}) ds \right) \middle| \mathcal{F}_\nu \right] \\ &= \mathbf{1}_A E_{Q_1} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_1}) ds \middle| \mathcal{F}_\nu \right] + \mathbf{1}_{A^c} E_{Q_2} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_2}) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.} \end{aligned} \quad (5.17)$$

Letting $A = \left\{ E_{Q_1} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_1}) ds \middle| \mathcal{F}_\nu \right] \leq E_{Q_2} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_2}) ds \middle| \mathcal{F}_\nu \right] \right\} \in \mathcal{F}_\nu$ above, one obtains that

$$E_{Q_3} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_3}) ds \middle| \mathcal{F}_\nu \right] = E_{Q_1} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_1}) ds \middle| \mathcal{F}_\nu \right] \wedge E_{Q_2} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_2}) ds \middle| \mathcal{F}_\nu \right] \quad P\text{-a.s.}$$

proving (5.13). Appealing to the basic properties of the essential infimum (e.g., Neveu [1975, Proposition VI-1-1]), we can find a sequence $\{Q_n^{\gamma, k}\}_{n \in \mathbb{N}}$ in $\mathcal{Q}_\nu^k$ such that (3.9) holds.

(2) Taking essential suprema over $\gamma \in \mathcal{S}_{\nu, T}$ on both sides of (5.17), we can deduce from Lemma 2.1 that

$$\begin{aligned} R^{Q_3}(\nu) &= \operatorname{ess\,sup}_{\gamma \in \mathcal{S}_{\nu, T}} E_{Q_3} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_3}) ds \middle| \mathcal{F}_\nu \right] \\ &= \mathbf{1}_A \operatorname{ess\,sup}_{\gamma \in \mathcal{S}_{\nu, T}} E_{Q_1} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_1}) ds \middle| \mathcal{F}_\nu \right] + \mathbf{1}_{A^c} \operatorname{ess\,sup}_{\gamma \in \mathcal{S}_{\nu, T}} E_{Q_2} \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^{Q_2}) ds \middle| \mathcal{F}_\nu \right] \\ &= \mathbf{1}_A R^{Q_1}(\nu) + \mathbf{1}_{A^c} R^{Q_2}(\nu), \quad P\text{-a.s.} \end{aligned}$$

Taking $A = \{R^{Q_1}(\nu) \leq R^{Q_2}(\nu)\} \in \mathcal{F}_\nu$ yields that $R^{Q_3}(\nu) = R^{Q_1}(\nu) \wedge R^{Q_2}(\nu)$, P -a.s., thus the family $\{R^Q(\nu)\}_{Q \in \mathcal{Q}_\nu^k}$ is directed downwards. Applying Proposition VI-1-1 of Neveu [1975] once again, one can find a sequence $\{Q_n^{(k)}\}_{n \in \mathbb{N}}$ in $\mathcal{Q}_\nu^k$ such that (3.10) holds. $\square$

Proof of Lemma 3.3: Let $Q_1, Q_2 \in \mathcal{Q}_\nu^k$. We define the stopping time $\gamma \triangleq \tau^{Q_1}(\nu) \wedge \tau^{Q_2}(\nu) \in \mathcal{S}_{\nu, T}$ and the event $A \triangleq \{R_\gamma^{Q_1, 0} \leq R_\gamma^{Q_2, 0}\} \in \mathcal{F}_\gamma$. It is clear that

$$\theta_t^{Q_3} \triangleq \mathbf{1}_{\{t > \gamma\}} \left(\mathbf{1}_A \theta_t^{Q_1} + \mathbf{1}_{A^c} \theta_t^{Q_2} \right), \quad t \in [0, T] \quad (5.18)$$

forms a predictable process, thus we can define a probability measure $Q_3 \in \mathcal{M}^e$ by $(dQ_3/dP) \triangleq \mathcal{E}(\theta^{Q_3} \bullet B)_T$. By analogy with (5.15), we have

$$f(t, \theta_t^{Q_3}) = \mathbf{1}_{\{t > \gamma\}} \left(\mathbf{1}_A f(t, \theta_t^{Q_1}) + \mathbf{1}_{A^c} f(t, \theta_t^{Q_2}) \right), \quad dt \otimes dP\text{-a.e.} \quad (5.19)$$

which together with (5.18) implies that $\theta^{Q_3} = 0$, $dt \otimes dP$ -a.e. on $\llbracket 0, \gamma \rrbracket$ and $|\theta_t^{Q_3}(\omega)| \vee f(t, \omega, \theta_t^{Q_3}(\omega)) \leq k$, $dt \otimes dP$ -a.e. on $\llbracket \gamma, T \rrbracket$. Hence $Q_3 \in \mathcal{Q}_\gamma^k \subset \mathcal{Q}_\nu^k$, thanks to Remark 3.2. Moreover, by analogy with (5.16), we can deduce that for any $\zeta \in \mathcal{S}_{\gamma, T}$ we have

$$Z_{\gamma, \zeta}^{Q_3} = \mathbf{1}_A Z_{\gamma, \zeta}^{Q_1} + \mathbf{1}_{A^c} Z_{\gamma, \zeta}^{Q_2}, \quad P\text{-a.s.} \quad (5.20)$$

Now fix $t \in [0, T]$. For any $\sigma \in \mathcal{S}_{\gamma \vee t, T}$, (5.20) shows that

$$Z_{\gamma \vee t, \sigma}^{Q_3} = \frac{Z_{\gamma, \sigma}^{Q_3}}{Z_{\gamma, \gamma \vee t}^{Q_3}} = \mathbf{1}_A \frac{Z_{\gamma, \sigma}^{Q_1}}{Z_{\gamma, \gamma \vee t}^{Q_1}} + \mathbf{1}_{A^c} \frac{Z_{\gamma, \sigma}^{Q_2}}{Z_{\gamma, \gamma \vee t}^{Q_2}} = \mathbf{1}_A Z_{\gamma \vee t, \sigma}^{Q_1} + \mathbf{1}_{A^c} Z_{\gamma \vee t, \sigma}^{Q_2}, \quad P\text{-a.s.},$$

and Bayes' Rule together with (5.19) imply then

$$\begin{aligned} E_{Q_3} \left[Y_\sigma + \int_{\gamma \vee t}^\sigma f(s, \theta_s^{Q_3}) ds \middle| \mathcal{F}_{\gamma \vee t} \right] &= E \left[Z_{\gamma \vee t, \sigma}^{Q_3} \left(Y_\sigma + \int_{\gamma \vee t}^\sigma f(s, \theta_s^{Q_3}) ds \right) \middle| \mathcal{F}_{\gamma \vee t} \right] \\ &= E \left[\mathbf{1}_A Z_{\gamma \vee t, \sigma}^{Q_1} \left(Y_\sigma + \int_{\gamma \vee t}^\sigma f(s, \theta_s^{Q_1}) ds \right) + \mathbf{1}_{A^c} Z_{\gamma \vee t, \sigma}^{Q_2} \left(Y_\sigma + \int_{\gamma \vee t}^\sigma f(s, \theta_s^{Q_2}) ds \right) \middle| \mathcal{F}_{\gamma \vee t} \right] \\ &= \mathbf{1}_A E_{Q_1} \left[Y_\sigma + \int_{\gamma \vee t}^\sigma f(s, \theta_s^{Q_1}) ds \middle| \mathcal{F}_{\gamma \vee t} \right] + \mathbf{1}_{A^c} E_{Q_2} \left[Y_\sigma + \int_{\gamma \vee t}^\sigma f(s, \theta_s^{Q_2}) ds \middle| \mathcal{F}_{\gamma \vee t} \right], \quad P\text{-a.s.} \end{aligned}$$

Taking essential suprema over $\sigma \in \mathcal{S}_{\gamma \vee t, T}$ on both sides above, we can deduce from Lemma 2.1 as well as (3.3) that

$$R_{\gamma \vee t}^{Q_3, 0} = R^{Q_3}(\gamma \vee t) = \mathbf{1}_A R^{Q_1}(\gamma \vee t) + \mathbf{1}_{A^c} R^{Q_2}(\gamma \vee t) = \mathbf{1}_A R_{\gamma \vee t}^{Q_1, 0} + \mathbf{1}_{A^c} R_{\gamma \vee t}^{Q_2, 0}, \quad P\text{-a.s.}$$

Since $R^{Q_i, 0}$, $i = 1, 2, 3$ are all RCLL processes, we have $R_{\gamma \vee t}^{Q_3, 0} = \mathbf{1}_A R_{\gamma \vee t}^{Q_1, 0} + \mathbf{1}_{A^c} R_{\gamma \vee t}^{Q_2, 0}$, $\forall t \in [0, T]$ outside a null set N , and this implies

$$\begin{aligned} \tau^{Q_3}(\nu) &= \inf \left\{ t \in [\nu, T] : R_t^{Q_3, 0} = Y_t \right\} \leq \inf \left\{ t \in [\gamma, T] : R_t^{Q_3, 0} = Y_t \right\} \\ &= \mathbf{1}_A \inf \left\{ t \in [\gamma, T] : R_t^{Q_1, 0} = Y_t \right\} + \mathbf{1}_{A^c} \inf \left\{ t \in [\gamma, T] : R_t^{Q_2, 0} = Y_t \right\}, \quad P\text{-a.s.} \end{aligned} \quad (5.21)$$

Since $R_{\tau^{Q_j}(\nu)}^{Q_j, 0} = Y_{\tau^{Q_j}(\nu)}$, P -a.s. for $j = 1, 2$, and since $\gamma = \tau^{Q_1}(\nu) \wedge \tau^{Q_2}(\nu)$, it holds P -a.s. that Y_γ is equal either to $R_\gamma^{Q_1, 0}$ or to $R_\gamma^{Q_2, 0}$. Then the definition of the set A shows that $R_\gamma^{Q_1, 0} = Y_\gamma$ holds P -a.s. on A , and that $R_\gamma^{Q_2, 0} = Y_\gamma$ holds P -a.s. on A^c , both of which further imply that

$$\mathbf{1}_A \inf \left\{ t \in [\gamma, T] : R_t^{Q_1, 0} = Y_t \right\} = \gamma \mathbf{1}_A \quad \text{and} \quad \mathbf{1}_{A^c} \inf \left\{ t \in [\gamma, T] : R_t^{Q_2, 0} = Y_t \right\} = \gamma \mathbf{1}_{A^c}, \quad P\text{-a.s.}$$

We conclude from (5.21) that $\tau^{Q_3}(\nu) \leq \gamma = \tau^{Q_1}(\nu) \wedge \tau^{Q_2}(\nu)$ holds P -a.s., hence the family $\{\tau^Q(\nu)\}_{Q \in \mathcal{Q}_\nu^k}$ is directed downwards. Thanks to Neveu [1975, page 121], we can find a sequence $\{Q_n^{(k)}\}_{n \in \mathbb{N}}$ in $\mathcal{Q}_\nu^k$, such that $\tau_k(\nu) = \text{ess inf}_{Q \in \mathcal{Q}_\nu^k} \tau^Q(\nu) = \lim_{n \rightarrow \infty} \downarrow \tau^{Q_n^{(k)}}(\nu)$, P -a.s. The limit $\lim_{n \rightarrow \infty} \downarrow \tau^{Q_n^{(k)}}(\nu)$ is also a stopping time in $\mathcal{S}_{\nu, T}$. $\square$

Proof of Lemma 3.4: It is easy to see from (3.12) and (f3) that

$$\theta^{Q'} = \theta^Q = 0, \quad dt \otimes dP\text{-a.e. on } \llbracket 0, \nu \rrbracket, \quad (5.22)$$

and that

$$f(t, \theta_t^{Q'}) = \mathbf{1}_{\{t \leq \gamma\}} f(t, \theta_t^Q) + \mathbf{1}_{\{t > \gamma\}} f(t, \tilde{\theta}_t^Q), \quad dt \otimes dP\text{-a.e.} \quad (5.23)$$

As a result

$$\begin{aligned} E_{Q'} \int_\nu^T f(s, \theta_s^{Q'}) ds &= E_{Q'} \int_\nu^\gamma f(s, \theta_s^Q) ds + E_{Q'} \int_\gamma^T f(s, \tilde{\theta}_s^Q) ds \\ &\leq E_Q \int_\nu^\gamma f(s, \theta_s^Q) ds + E_{Q'} \int_\gamma^T k ds \leq E_Q \int_\nu^T f(s, \theta_s^Q) ds + kT < \infty, \end{aligned}$$

thus $Q' \in \mathcal{Q}_\nu$. If $Q \in \mathcal{Q}_\nu^k$, we see from (3.12) and (5.23) that

$$|\theta_t^{Q'}(\omega)| \vee f(t, \omega, \theta_t^{Q'}(\omega)) = \begin{cases} |\theta_t^Q(\omega)| \vee f(t, \omega, \theta_t^Q(\omega)) \leq k & dt \otimes dP\text{-a.e. on } \llbracket \nu, \gamma \rrbracket, \\ |\tilde{\theta}_t^Q(\omega)| \vee f(t, \omega, \tilde{\theta}_t^Q(\omega)) \leq k & dt \otimes dP\text{-a.e. on } \llbracket \gamma, T \rrbracket, \end{cases}$$

which, together with (5.22), shows that $Q' \in \mathcal{Q}_\nu^k$.

Now we fix $\sigma \in \mathcal{S}_{\gamma, T}$. For any $\delta \in \mathcal{S}_{\sigma, T}$, Bayes' Rule shows

$$E_{Q'} \left[Y_\delta + \int_\sigma^\delta f(s, \theta_s^{Q'}) ds \middle| \mathcal{F}_\sigma \right] = E_{Q'} \left[Y_\delta + \int_\sigma^\delta f(s, \theta_s^{\tilde{Q}}) ds \middle| \mathcal{F}_\sigma \right] = E_{\tilde{Q}} \left[Y_\delta + \int_\sigma^\delta f(s, \theta_s^{\tilde{Q}}) ds \middle| \mathcal{F}_\sigma \right], \quad P\text{-a.s.},$$

and (3.3) implies

$$\begin{aligned} R_\sigma^{Q', 0} = R^{Q'}(\sigma) &= \operatorname{ess\,sup}_{\delta \in \mathcal{S}_{\sigma, T}} E_{Q'} \left[Y_\delta + \int_\sigma^\delta f(s, \theta_s^{Q'}) ds \middle| \mathcal{F}_\sigma \right] \\ &= \operatorname{ess\,sup}_{\delta \in \mathcal{S}_{\sigma, T}} E_{\tilde{Q}} \left[Y_\delta + \int_\sigma^\delta f(s, \theta_s^{\tilde{Q}}) ds \middle| \mathcal{F}_\sigma \right] = R^{\tilde{Q}}(\sigma) = R_\sigma^{\tilde{Q}, 0}, \quad P\text{-a.s.} \end{aligned} \quad \square$$

Proof of Theorem 3.1: Fix $Q \in \mathcal{Q}_\nu$. For any $m, k \in \mathbb{N}$, we consider the probability measure $Q^{m, k} \in \mathcal{Q}_\nu^k$ as defined in (5.5). In light of Lemma 3.3, for any $l \in \mathbb{N}$ there exists a sequence $\{Q_n^{(l)}\}_{n \in \mathbb{N}}$ in $\mathcal{Q}_\nu^l$ such that $\tau_l(\nu) = \lim_{n \rightarrow \infty} \downarrow \tau_{Q_n^{(l)}}(\nu)$, P -a.s. Now let $k, l, m, n \in \mathbb{N}$ with $k \leq l$. Lemma 3.4 implies that the predictable process

$$\theta_t^{Q_n^{m, k, l}} \triangleq \mathbf{1}_{\{t \leq \tau_l(\nu)\}} \theta_t^{Q_n^{m, k}} + \mathbf{1}_{\{t > \tau_l(\nu)\}} \theta_t^{Q_n^{(l)}}, \quad t \in [0, T]$$

induces a probability measure $Q_n^{m, k, l} \in \mathcal{Q}_\nu^l$ via $dQ_n^{m, k, l} = \mathcal{E}(\theta^{Q_n^{m, k, l}} \bullet B)_T dP$, such that for any $t \in [0, T]$, we have $R_{\tau_l(\nu) \vee t}^{Q_n^{m, k, l}, 0} = R_{\tau_l(\nu) \vee t}^{Q_n^{(l)}, 0}$, P -a.s. Since $R_{\tau_l(\nu) \vee t}^{Q_n^{m, k, l}, 0}$ and $R_{\tau_l(\nu) \vee t}^{Q_n^{(l)}, 0}$ are both RCLL processes, outside a null set N we have

$$R_{\tau_l(\nu) \vee t}^{Q_n^{m, k, l}, 0} = R_{\tau_l(\nu) \vee t}^{Q_n^{(l)}, 0}, \quad \forall t \in [0, T]$$

and this, together with the fact that $\tau_l(\nu) \leq \tau_{Q_n^{m, k, l}}(\nu) \wedge \tau_{Q_n^{(l)}}(\nu)$, P -a.s. implies

$$\begin{aligned} \tau_{Q_n^{m, k, l}}(\nu) &= \inf \left\{ t \in [\nu, T] : R_t^{Q_n^{m, k, l}, 0} = Y_t \right\} = \inf \left\{ t \in [\tau_l(\nu), T] : R_t^{Q_n^{m, k, l}, 0} = Y_t \right\} \\ &= \inf \left\{ t \in [\tau_l(\nu), T] : R_t^{Q_n^{(l)}, 0} = Y_t \right\} = \inf \left\{ t \in [\nu, T] : R_t^{Q_n^{(l)}, 0} = Y_t \right\} = \tau_{Q_n^{(l)}}(\nu), \quad P\text{-a.s.} \end{aligned} \quad (5.24)$$

Similar to (5.6), we have

$$f(t, \theta_t^{Q_n^{m, k, l}}) = \mathbf{1}_{\{t \leq \tau_l(\nu)\}} f(t, \theta_t^{Q_n^{m, k}}) + \mathbf{1}_{\{t > \tau_l(\nu)\}} f(t, \theta_t^{Q_n^{(l)}}), \quad dt \otimes dP\text{-a.e.} \quad (5.25)$$

Then one can deduce from (5.24) and (5.25) that

$$\begin{aligned} \bar{V}(\nu) &= \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} R^Q(\nu) \leq R^{Q_n^{m, k, l}}(\nu) = E_{Q_n^{m, k, l}} \left(Y_{\tau_{Q_n^{m, k, l}}(\nu)} + \int_\nu^{\tau_{Q_n^{m, k, l}}(\nu)} f(s, \theta_s^{Q_n^{m, k, l}}) ds \middle| \mathcal{F}_\nu \right) \\ &= E_{Q_n^{m, k, l}} \left[Y_{\tau_{Q_n^{(l)}}(\nu)} + \int_{\tau_l(\nu)}^{\tau_{Q_n^{(l)}}(\nu)} f(s, \theta_s^{Q_n^{m, k, l}}) ds \middle| \mathcal{F}_\nu \right] + E_{Q_n^{m, k}} \left[\int_\nu^{\tau_l(\nu)} f(s, \theta_s^{Q_n^{m, k, l}}) ds \middle| \mathcal{F}_\nu \right] \\ &= E \left[\left(Z_{\nu, \tau_{Q_n^{(l)}}(\nu)}^{Q_n^{m, k, l}} - Z_{\nu, \tau_l(\nu)}^{Q_n^{m, k}} \right) \cdot \left(Y_{\tau_{Q_n^{(l)}}(\nu)} + \int_{\tau_l(\nu)}^{\tau_{Q_n^{(l)}}(\nu)} f(s, \theta_s^{Q_n^{(l)}}) ds \right) \middle| \mathcal{F}_\nu \right] \\ &\quad + E \left[Z_{\nu, \tau_l(\nu)}^{Q_n^{m, k}} \left(Y_{\tau_{Q_n^{(l)}}(\nu)} + \int_{\tau_l(\nu)}^{\tau_{Q_n^{(l)}}(\nu)} f(s, \theta_s^{Q_n^{(l)}}) ds \right) \middle| \mathcal{F}_\nu \right] + E_{Q_n^{m, k}} \left[\int_\nu^{\tau_l(\nu)} f(s, \theta_s^{Q_n^{m, k}}) ds \middle| \mathcal{F}_\nu \right] \\ &\leq (\|Y\|_\infty + lT) \cdot E \left[\left| Z_{\nu, \tau_{Q_n^{(l)}}(\nu)}^{Q_n^{m, k, l}} - Z_{\nu, \tau_l(\nu)}^{Q_n^{m, k}} \right| \middle| \mathcal{F}_\nu \right] + E \left[Z_{\nu, \tau_l(\nu)}^{Q_n^{m, k}} \left(Y_{\tau_{Q_n^{(l)}}(\nu)} + k(\tau_{Q_n^{(l)}}(\nu) - \tau_l(\nu)) \right) \middle| \mathcal{F}_\nu \right] \\ &\quad + E_{Q_n^{m, k}} \left[\int_\nu^{\tau_l(\nu)} f(s, \theta_s^{Q_n^{m, k}}) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.} \end{aligned} \quad (5.26)$$

Because $E\left(\int_{\tau_l(\nu)}^{\tau^{Q_n^{(l)}}(\nu)} \theta_s^{Q_n^{(l)}} dB_s\right)^2 = E\int_{\tau_l(\nu)}^{\tau^{Q_n^{(l)}}(\nu)} |\theta_s^{Q_n^{(l)}}|^2 ds \leq l^2 E\left[\tau^{Q_n^{(l)}}(\nu) - \tau_l(\nu)\right]$, which goes to zero as $n \rightarrow \infty$, using similar arguments to those that lead to (5.8), we can find a subsequence of $\{Q_n^{(l)}\}_{n \in \mathbb{N}}$ (we still denote it by $\{Q_n^{(l)}\}_{n \in \mathbb{N}}$) such that $\lim_{n \rightarrow \infty} Z_{\nu, \tau^{Q_n^{(l)}}(\nu)}^{Q_n^{m,k,l}} = Z_{\nu, \tau_l(\nu)}^{Q^{m,k}}$, P -a.s. Since $E\left[Z_{\nu, \tau^{Q_n^{(l)}}(\nu)}^{Q_n^{m,k,l}} \middle| \mathcal{F}_\nu\right] = E\left[Z_{\nu, \tau_l(\nu)}^{Q^{m,k}} \middle| \mathcal{F}_\nu\right] = 1$, P -a.s. for any $n \in \mathbb{N}$, Scheffé's Lemma implies

$$\lim_{n \rightarrow \infty} E\left(\left|Z_{\nu, \tau^{Q_n^{(l)}}(\nu)}^{Q_n^{m,k,l}} - Z_{\nu, \tau_l(\nu)}^{Q^{m,k}}\right| \middle| \mathcal{F}_\nu\right) = 0, \quad P\text{-a.s.} \quad (5.27)$$

On the other hand, since

$$Z_{\nu, \tau_l(\nu)}^{Q^{m,k}} \left| Y_{\tau^{Q_n^{(l)}}(\nu)} + k(\tau^{Q_n^{(l)}}(\nu) - \tau_l(\nu)) \right| \leq Z_{\nu, \tau_l(\nu)}^{Q^{m,k}} (\|Y\|_\infty + kT), \quad P\text{-a.s.},$$

and since Y is right-continuous, the Dominated Convergence Theorem gives

$$\lim_{n \rightarrow \infty} E\left[Z_{\nu, \tau_l(\nu)}^{Q^{m,k}} \left(Y_{\tau^{Q_n^{(l)}}(\nu)} + k(\tau^{Q_n^{(l)}}(\nu) - \tau_l(\nu))\right) \middle| \mathcal{F}_\nu\right] = E\left[Z_{\nu, \tau_l(\nu)}^{Q^{m,k}} Y_{\tau_l(\nu)} \middle| \mathcal{F}_\nu\right] = E_{Q^{m,k}}[Y_{\tau_l(\nu)} | \mathcal{F}_\nu], \quad P\text{-a.s.} \quad (5.28)$$

Therefore, letting $n \rightarrow \infty$ in (5.26), we can deduce from (5.27) and (5.28) that

$$\bar{V}(\nu) \leq E_{Q^{m,k}}\left[Y_{\tau_l(\nu)} + \int_{\nu}^{\tau_l(\nu)} f(s, \theta_s^{Q^{m,k}}) ds \middle| \mathcal{F}_\nu\right], \quad P\text{-a.s.}$$

As $l \rightarrow \infty$, the Bounded Convergence Theorem gives

$$\bar{V}(\nu) \leq E_{Q^{m,k}}\left[Y_{\tau(\nu)} + \int_{\nu}^{\tau(\nu)} f(s, \theta_s^{Q^{m,k}}) ds \middle| \mathcal{F}_\nu\right], \quad P\text{-a.s.}$$

whence, just as in (5.7), we deduce

$$\begin{aligned} \bar{V}(\nu) &\leq E_{Q^{m,k}}\left[Y_{\tau(\nu)} + \int_{\nu}^{\tau(\nu)} f(s, \theta_s^{Q^{m,k}}) ds \middle| \mathcal{F}_\nu\right] \\ &\leq (\|Y\|_\infty + m) \cdot E\left[\left|Z_{\nu, \tau(\nu)}^{Q^{m,k}} - Z_{\nu, \tau(\nu) \wedge \delta_m^Q}^Q\right| \middle| \mathcal{F}_\nu\right] + \|Y\|_\infty \cdot E\left[\left|Z_{\nu, \tau(\nu) \wedge \delta_m^Q}^Q - Z_{\nu, \tau(\nu)}^Q\right| \middle| \mathcal{F}_\nu\right] \\ &\quad + E_Q\left[Y_{\tau(\nu)} + \int_{\nu}^{\tau(\nu)} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu\right], \quad P\text{-a.s.} \end{aligned} \quad (5.29)$$

By analogy with (5.8) and (5.10), one can show that for any $m \in \mathbb{N}$ we have $\lim_{k \rightarrow \infty} E\left[\left|Z_{\nu, \tau(\nu)}^{Q^{m,k}} - Z_{\nu, \tau(\nu) \wedge \delta_m^Q}^Q\right| \middle| \mathcal{F}_\nu\right] = 0$, P -a.s. and that $\lim_{m \rightarrow \infty} E\left[\left|Z_{\nu, \tau(\nu) \wedge \delta_m^Q}^Q - Z_{\nu, \tau(\nu)}^Q\right| \middle| \mathcal{F}_\nu\right] = 0$, P -a.s. Therefore, letting $k \rightarrow \infty$ and subsequently letting $m \rightarrow \infty$ in (5.29), we obtain

$$\bar{V}(\nu) \leq E_Q\left[Y_{\tau(\nu)} + \int_{\nu}^{\tau(\nu)} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu\right], \quad P\text{-a.s.}$$

Taking the essential infimum of the right-hand-side over $Q \in \mathcal{Q}_\nu$ yields

$$\begin{aligned} \bar{V}(\nu) &\leq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q\left[Y_{\tau(\nu)} + \int_{\nu}^{\tau(\nu)} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu\right] \\ &\leq \operatorname{ess\,sup}_{\gamma \in \mathcal{S}_{\nu, T}} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q\left[Y_\gamma + \int_{\nu}^{\gamma} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu\right] = \underline{V}(\nu) \leq \bar{V}(\nu), \quad P\text{-a.s.} \end{aligned}$$

and the result follows. $\square$

Proof of Proposition 3.2: For each fixed $k \in \mathbb{N}$, there exists on the strength of Lemma 3.3 a sequence $\{Q_n^{(k)}\}_{n \in \mathbb{N}}$ in $\mathcal{Q}_\nu^k$ such that $\tau_k(\nu) = \lim_{n \rightarrow \infty} \downarrow \tau^{Q_n^{(k)}}(\nu)$, P -a.s.

For any $n \in \mathbb{N}$, the predictable process $\theta_t^{\tilde{Q}_n^{(k)}} \triangleq \mathbf{1}_{\{t > \tau_k(\nu)\}} \theta_t^{Q_n^{(k)}}$, $t \in [0, T]$ induces a probability measure $\tilde{Q}_n^{(k)}$ by $d\tilde{Q}_n^{(k)} \triangleq \mathcal{E}(\tilde{Q}_n^{(k)} \bullet B)_T dP = Z_{\tau_k(\nu), T}^{Q_n^{(k)}} dP$. Since $\nu \leq \sigma \triangleq \tau(\nu) \leq \tau_k(\nu) \leq \tau^{\tilde{Q}_n^{(k)}}(\nu)$, P -a.s., we have $\tilde{Q}_n^{(k)} \in \mathcal{Q}_{\tau_k(\nu)}^k \subset \mathcal{Q}_\sigma^k \subset \mathcal{Q}_\nu^k$ and

$$\tau^{\tilde{Q}_n^{(k)}}(\nu) = \inf \{t \in [\nu, T] : R_t^{\tilde{Q}_n^{(k)}, 0} = Y_t\} = \inf \{t \in [\sigma, T] : R_t^{\tilde{Q}_n^{(k)}, 0} = Y_t\} = \tau^{\tilde{Q}_n^{(k)}}(\sigma), \quad P\text{-a.s.} \quad (5.30)$$

We also know from Lemma 3.4 that for any $t \in [0, T]$: $R_{\tau_k(\nu) \vee t}^{\tilde{Q}_n^{(k)}, 0} = R_{\tau_k(\nu) \vee t}^{Q_n^{(k)}, 0}$, P -a.s.

Since $R_{\tau_k(\nu), 0}^{\tilde{Q}_n^{(k)}}$ and $R_{\tau_k(\nu), 0}^{Q_n^{(k)}}$ are both RCLL processes, there exists a null set N outside which we have $R_{\tau_k(\nu) \vee t}^{\tilde{Q}_n^{(k)}, 0} = R_{\tau_k(\nu) \vee t}^{Q_n^{(k)}, 0}$, $\forall t \in [0, T]$. By analogy with (5.24) and (5.6), respectively, we have

$$\tau^{\tilde{Q}_n^{(k)}}(\nu) = \tau^{Q_n^{(k)}}(\nu), \quad P\text{-a.s.} \quad (5.31)$$

and $f(t, \theta_t^{\tilde{Q}_n^{(k)}}) = \mathbf{1}_{\{t > \tau_k(\nu)\}} f(t, \theta_t^{Q_n^{(k)}})$, $dt \otimes dP$ -a.e. Then we can deduce from (5.30), (5.31) that

$$\begin{aligned} V(\sigma) &= \bar{V}(\sigma) = \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\sigma} R^Q(\sigma) \leq R^{\tilde{Q}_n^{(k)}}(\sigma) = E_{\tilde{Q}_n^{(k)}} \left(Y_{\tau^{Q_n^{(k)}}(\nu)} + \int_{\sigma}^{\tau^{Q_n^{(k)}}(\nu)} \mathbf{1}_{\{s > \tau_k(\nu)\}} f(s, \theta_s^{Q_n^{(k)}}) ds \middle| \mathcal{F}_\sigma \right) \\ &= E_{\tilde{Q}_n^{(k)}} \left[Y_{\tau^{Q_n^{(k)}}(\nu)} + \int_{\sigma}^{\tau^{Q_n^{(k)}}(\nu)} \mathbf{1}_{\{s > \tau_k(\nu)\}} f(s, \theta_s^{Q_n^{(k)}}) ds \middle| \mathcal{F}_\sigma \right] \\ &= E \left[\left(Z_{\sigma, \tau^{Q_n^{(k)}}(\nu)}^{\tilde{Q}_n^{(k)}} - 1 \right) \cdot \left(Y_{\tau^{Q_n^{(k)}}(\nu)} + \int_{\tau_k(\nu)}^{\tau^{Q_n^{(k)}}(\nu)} f(s, \theta_s^{Q_n^{(k)}}) ds \right) \middle| \mathcal{F}_\sigma \right] \\ &\quad + E \left[Y_{\tau^{Q_n^{(k)}}(\nu)} + \int_{\tau_k(\nu)}^{\tau^{Q_n^{(k)}}(\nu)} f(s, \theta_s^{Q_n^{(k)}}) ds \middle| \mathcal{F}_\sigma \right] \\ &\leq (\|Y\|_\infty + kT) \cdot E \left[\left| Z_{\tau_k(\nu), \tau^{Q_n^{(k)}}(\nu)}^{Q_n^{(k)}} - 1 \right| \middle| \mathcal{F}_\sigma \right] + E \left[Y_{\tau^{Q_n^{(k)}}(\nu)} + k(\tau^{Q_n^{(k)}}(\nu) - \tau_k(\nu)) \middle| \mathcal{F}_\sigma \right], \quad P\text{-a.s.} \quad (5.32) \end{aligned}$$

Just as in (5.27), it can shown that $\lim_{n \rightarrow \infty} E \left(\left| Z_{\tau_k(\nu), \tau^{Q_n^{(k)}}(\nu)}^{Q_n^{(k)}} - 1 \right| \middle| \mathcal{F}_\sigma \right) = 0$, P -a.s.; on the other hand, the Bounded Convergence Theorem implies

$$\lim_{n \rightarrow \infty} E \left(Y_{\tau^{Q_n^{(k)}}(\nu)} + k(\tau^{Q_n^{(k)}}(\nu) - \tau_k(\nu)) \middle| \mathcal{F}_\sigma \right) = E \left[Y_{\tau_k(\nu)} \middle| \mathcal{F}_\sigma \right], \quad P\text{-a.s.}$$

Letting $n \rightarrow \infty$ in (5.32) yields $V(\sigma) \leq E \left[Y_{\tau_k(\nu)} \middle| \mathcal{F}_\sigma \right]$, P -a.s., and applying the Bounded Convergence Theorem we obtain $V(\sigma) \leq \lim_{k \rightarrow \infty} E \left[Y_{\tau_k(\nu)} \middle| \mathcal{F}_\sigma \right] = E \left[Y_\sigma \middle| \mathcal{F}_\sigma \right] = Y_\sigma$, P -a.s. The reverse inequality is rather obvious. $\square$

Proof of Proposition 3.3: Fix $k \in \mathbb{N}$. In light of (3.10), we can find a sequence $\{Q_n^{(k)}\}_{n \in \mathbb{N}} \subset \mathcal{Q}_\gamma^k$ such that

$$\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\gamma^k} R^Q(\gamma) = \lim_{n \rightarrow \infty} \downarrow R^{Q_n^{(k)}}(\gamma), \quad P\text{-a.s.} \quad (5.33)$$

For any $n \in \mathbb{N}$, Lemma 3.4 implies that the predictable process $\theta_t^{\tilde{Q}_n^{(k)}} \triangleq \mathbf{1}_{\{t \leq \gamma\}} \theta_t^Q + \mathbf{1}_{\{t > \gamma\}} \theta_t^{Q_n^{(k)}}$, $t \in [0, T]$ induces a probability measure $\tilde{Q}_n^{(k)} \in \mathcal{Q}_\gamma$ via $d\tilde{Q}_n^{(k)} \triangleq \mathcal{E}(\theta^{\tilde{Q}_n^{(k)}} \bullet B)_T dP$, such that for any $t \in [0, T]$, $R_t^{\tilde{Q}_n^{(k)}}(\gamma) = R_t^{Q_n^{(k)}}(\gamma)$, P -a.s. Since $\gamma \leq \tau(\nu) \leq \tau^{\tilde{Q}_n^{(k)}}(\nu)$, P -a.s., applying (3.4) yields

$$\begin{aligned} V(\nu) &\leq R^{\tilde{Q}_n^{(k)}}(\nu) = E_{\tilde{Q}_n^{(k)}} \left[R^{\tilde{Q}_n^{(k)}}(\gamma) + \int_\nu^\gamma f(s, \theta_s^{\tilde{Q}_n^{(k)}}) ds \middle| \mathcal{F}_\nu \right] = E_{\tilde{Q}_n^{(k)}} \left[R^{Q_n^{(k)}}(\gamma) + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \\ &= E_Q \left[R^{Q_n^{(k)}}(\gamma) + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.} \quad (5.34) \end{aligned}$$

It follows from (3.2) that

$$-\|Y\|_\infty \leq Y_\gamma \leq R^{Q_n^{(k)}}(\gamma) \leq \|Y\|_\infty + kT, \quad P\text{-a.s.} \quad (5.35)$$

Letting $n \rightarrow \infty$ in (5.34), we can deduce from the Bounded Convergence Theorem that

$$V(\nu) \leq E_Q \left[\lim_{n \rightarrow \infty} \downarrow R^{Q_n^{(k)}}(\gamma) \middle| \mathcal{F}_\nu \right] + E_Q \left[\int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] = E_Q \left[\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\gamma^k} R^Q(\gamma) + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.}$$

Letting $n \rightarrow \infty$ in (5.35), one sees from (5.33) that $-\|Y\|_\infty \leq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\gamma^k} R^Q(\gamma) \leq \|Y\|_\infty + kT$ holds P -a.s., and this leads to

$$-\|Y\|_\infty \leq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\gamma^k} R^Q(\gamma) \leq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\gamma^1} R^Q(\gamma) \leq \|Y\|_\infty + T, \quad P\text{-a.s.}$$

From the Bounded Convergence Theorem and Lemma 3.1 we obtain now

$$V(\nu) \leq E_Q \left[\lim_{k \rightarrow \infty} \downarrow \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\gamma^k} R^Q(\gamma) \middle| \mathcal{F}_\nu \right] + E_Q \left[\int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] = E_Q \left[V(\gamma) + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.} \quad \square$$

Proof of Lemma 3.5: Fix $k \in \mathbb{N}$. For any $Q \in \mathcal{Q}_\nu^k$, the predictable process $\theta_t^{\tilde{Q}} \triangleq \mathbf{1}_{\{t > \nu \vee \gamma\}} \theta_t^Q$, $t \in [0, T]$ induces a probability measure $\tilde{Q}$ by $(d\tilde{Q}/dP) \triangleq \mathcal{E}(\tilde{Q} \bullet B)_T = Z_{\nu \vee \gamma, T}^Q$. Remark 3.2 shows that $\tilde{Q} \in \mathcal{Q}_{\nu \vee \gamma}^k \subset \mathcal{Q}_\nu^k \cap \mathcal{Q}_\gamma^k$. By analogy with (5.6), we have $f(t, \theta_t^{\tilde{Q}}) = \mathbf{1}_{\{t > \nu \vee \gamma\}} f(t, \theta_t^Q)$, $dt \otimes dP$ -a.e. Then one can deduce that

$$\begin{aligned} \mathbf{1}_{\{\nu=\gamma\}} E_{\tilde{Q}} \left[Y_{\sigma \vee \gamma} + \int_\gamma^{\sigma \vee \gamma} f(s, \theta_s^{\tilde{Q}}) ds \middle| \mathcal{F}_\gamma \right] &= \mathbf{1}_{\{\nu=\gamma\}} E_{\tilde{Q}} \left[Y_{\sigma \vee \gamma} + \int_\gamma^{\sigma \vee \gamma} \mathbf{1}_{\{s > \nu \vee \gamma\}} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \\ &= E_{\tilde{Q}} \left[\mathbf{1}_{\{\nu=\gamma\}} \left(Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \right) \middle| \mathcal{F}_\nu \right] = E \left[E_Q \left[\mathbf{1}_{\{\nu=\gamma\}} \left(Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \right) \middle| \mathcal{F}_{\nu \vee \gamma} \right] \middle| \mathcal{F}_\nu \right] \\ &= E \left[\mathbf{1}_{\{\nu=\gamma\}} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \middle| \mathcal{F}_\nu \right] = \mathbf{1}_{\{\nu=\gamma\}} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.}, \end{aligned} \quad (5.36)$$

which implies

$$\mathbf{1}_{\{\nu=\gamma\}} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \geq \mathbf{1}_{\{\nu=\gamma\}} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} E_Q \left[Y_{\sigma \vee \gamma} + \int_\gamma^{\sigma \vee \gamma} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\gamma \right], \quad P\text{-a.s.}$$

Taking the essential infimum of the left-hand-side over $Q \in \mathcal{Q}_\nu^k$, one can deduce from Lemma 2.1 that

$$\begin{aligned} \mathbf{1}_{\{\nu=\gamma\}} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] &= \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} \mathbf{1}_{\{\nu=\gamma\}} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \\ &\geq \mathbf{1}_{\{\nu=\gamma\}} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\gamma^k} E_Q \left[Y_{\sigma \vee \gamma} + \int_\gamma^{\sigma \vee \gamma} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\gamma \right], \quad P\text{-a.s.} \end{aligned}$$

Letting $k \rightarrow \infty$, we see from Lemma 3.1 (1) that

$$\mathbf{1}_{\{\nu=\gamma\}} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \geq \mathbf{1}_{\{\nu=\gamma\}} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\gamma} E_Q \left[Y_{\sigma \vee \gamma} + \int_\gamma^{\sigma \vee \gamma} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\gamma \right], \quad P\text{-a.s.}$$

Reversing the roles of ν and γ , we obtain (3.16).

On the other hand, taking essential supremum over $\sigma \in \mathcal{S}_{0,T}$ on both sides of (5.36), we can deduce from Lemma 2.1 that

$$\begin{aligned} \mathbf{1}_{\{\nu=\gamma\}} R^{\tilde{Q}}(\gamma) &= \operatorname{ess\,sup}_{\sigma \in \mathcal{S}_{0,T}} \mathbf{1}_{\{\nu=\gamma\}} E_{\tilde{Q}} \left[Y_{\sigma \vee \gamma} + \int_\gamma^{\sigma \vee \gamma} f(s, \theta_s^{\tilde{Q}}) ds \middle| \mathcal{F}_\gamma \right] \\ &= \operatorname{ess\,sup}_{\sigma \in \mathcal{S}_{0,T}} \mathbf{1}_{\{\nu=\gamma\}} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] = \mathbf{1}_{\{\nu=\gamma\}} R^Q(\nu), \quad P\text{-a.s.} \end{aligned}$$

which implies that $\mathbf{1}_{\{\nu=\gamma\}}R^Q(\nu) \geq \mathbf{1}_{\{\nu=\gamma\}}\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\gamma^k} R^Q(\gamma)$, P -a.s. Taking the essential infimum of the left-hand-side over $Q \in \mathcal{Q}_\nu^k$, one can deduce from Lemma 2.1 that

$$\mathbf{1}_{\{\nu=\gamma\}}\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} R^Q(\nu) = \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} \mathbf{1}_{\{\nu=\gamma\}} R^Q(\nu) \geq \mathbf{1}_{\{\nu=\gamma\}}\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\gamma^k} R^Q(\gamma), \quad P\text{-a.s.}$$

Letting $k \rightarrow \infty$, we see from Lemma 3.1 (2) that

$$\mathbf{1}_{\{\nu=\gamma\}}V(\nu) = \mathbf{1}_{\{\nu=\gamma\}}\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} R^Q(\nu) \geq \mathbf{1}_{\{\nu=\gamma\}}\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\gamma} R^Q(\gamma) = \mathbf{1}_{\{\nu=\gamma\}}V(\gamma), \quad P\text{-a.s.}$$

Reversing the roles of ν and γ , we obtain (3.17). $\square$

Proof of Theorem 3.2: Proof of (1).

Step 1: For any $\sigma, \nu \in \mathcal{S}_{0,T}$, we define

$$\Psi^\sigma(\nu) \triangleq \mathbf{1}_{\{\sigma \leq \nu\}}Y_\sigma + \mathbf{1}_{\{\sigma > \nu\}}\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right].$$

We see from (3.7) that

$$\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] = \lim_{k \rightarrow \infty} \downarrow \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.} \quad (5.37)$$

Fix $k \in \mathbb{N}$. In light of (3.9), we can find a sequence $\{Q_n^{(k)}\}_{n \in \mathbb{N}}$ in $\mathcal{Q}_\nu^k$ such that

$$\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] = \lim_{n \rightarrow \infty} \downarrow E_{Q_n^{(k)}} \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^{Q_n^{(k)}}) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.} \quad (5.38)$$

By analogy with (5.35), we have

$$- \|Y\|_\infty \leq E_{Q_n^{(k)}} \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^{Q_n^{(k)}}) ds \middle| \mathcal{F}_\nu \right] \leq \|Y\|_\infty + kT \quad (5.39)$$

P -a.s.; letting $n \rightarrow \infty$, we see from (5.38) that

$$- \|Y\|_\infty \leq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \leq \|Y\|_\infty + kT, \quad P\text{-a.s.}$$

Therefore,

$$\begin{aligned} - \|Y\|_\infty &\leq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \\ &\leq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^1} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \leq \|Y\|_\infty + T, \quad P\text{-a.s.} \end{aligned} \quad (5.40)$$

Letting $k \rightarrow \infty$, we see from (5.37) that

$$- \|Y\|_\infty \leq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \leq \|Y\|_\infty + T, \quad P\text{-a.s.}$$

which implies that

$$- \|Y\|_\infty \leq \Psi^\sigma(\nu) \leq \|Y\|_\infty + T, \quad P\text{-a.s.} \quad (5.41)$$

Let $\gamma \in \mathcal{S}_{0,T}$. It follows from (3.16) that

$$\begin{aligned} \mathbf{1}_{\{\nu=\gamma\}}\Psi^\sigma(\nu) &= \mathbf{1}_{\{\sigma \leq \nu=\gamma\}}Y_\sigma + \mathbf{1}_{\{\sigma > \nu=\gamma\}}\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \\ &= \mathbf{1}_{\{\sigma \leq \gamma=\nu\}}Y_\sigma + \mathbf{1}_{\{\sigma > \gamma=\nu\}}\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\gamma} E_Q \left[Y_{\sigma \vee \gamma} + \int_\gamma^{\sigma \vee \gamma} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\gamma \right] = \mathbf{1}_{\{\nu=\gamma\}}\Psi^\sigma(\gamma), \quad P\text{-a.s.} \end{aligned} \quad (5.42)$$

Step 2: Fix $\sigma \in \mathcal{S}_{0,T}$. For any $\zeta \in \mathcal{S}_{0,T}$, $\nu \in \mathcal{S}_{\zeta,T}$ and $k \in \mathbb{N}$, we let $\{Q_n^{(k)}\}_{n \in \mathbb{N}} \subset \mathcal{Q}_\nu^k$ be the sequence described in (5.38). Then we can deduce that

$$\begin{aligned} \Psi^\sigma(\zeta) &\leq \mathbf{1}_{\{\sigma \leq \zeta\}} Y_\sigma + \mathbf{1}_{\{\sigma > \zeta\}} E_{Q_n^{(k)}} \left[Y_{\sigma \vee \zeta} + \int_\zeta^{\sigma \vee \zeta} f(s, \theta_s^{Q_n^{(k)}}) ds \middle| \mathcal{F}_\zeta \right] \\ &= \mathbf{1}_{\{\sigma \leq \zeta\}} Y_{\sigma \wedge \zeta} + \mathbf{1}_{\{\sigma > \zeta\}} E \left[E_{Q_n^{(k)}} \left[Y_{\sigma \vee \zeta} + \int_\zeta^{\sigma \vee \zeta} f(s, \theta_s^{Q_n^{(k)}}) ds \middle| \mathcal{F}_\nu \right] \middle| \mathcal{F}_\zeta \right] \\ &= E \left[\mathbf{1}_{\{\sigma \leq \zeta\}} Y_{\sigma \wedge \zeta} + \mathbf{1}_{\{\sigma > \zeta\}} E_{Q_n^{(k)}} \left[Y_{\sigma \vee \zeta} + \int_\zeta^{\sigma \vee \zeta} f(s, \theta_s^{Q_n^{(k)}}) ds \middle| \mathcal{F}_\nu \right] \middle| \mathcal{F}_\zeta \right], \quad P\text{-a.s.} \end{aligned} \quad (5.43)$$

On the other hand, it holds P -a.s. that

$$\begin{aligned} \mathbf{1}_{\{\sigma > \nu\}} E_{Q_n^{(k)}} \left[Y_{\sigma \vee \zeta} + \int_\zeta^{\sigma \vee \zeta} f(s, \theta_s^{Q_n^{(k)}}) ds \middle| \mathcal{F}_\nu \right] &= E_{Q_n^{(k)}} \left[\mathbf{1}_{\{\sigma > \nu\}} \left(Y_\sigma + \int_\zeta^\sigma f(s, \theta_s^{Q_n^{(k)}}) ds \right) \middle| \mathcal{F}_\nu \right] \\ &= E_{Q_n^{(k)}} \left[\mathbf{1}_{\{\sigma > \nu\}} \left(Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^{Q_n^{(k)}}) ds \right) \middle| \mathcal{F}_\nu \right] = \mathbf{1}_{\{\sigma > \nu\}} E_{Q_n^{(k)}} \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^{Q_n^{(k)}}) ds \middle| \mathcal{F}_\nu \right] \end{aligned}$$

and that

$$\begin{aligned} \mathbf{1}_{\{\zeta < \sigma \leq \nu\}} E_{Q_n^{(k)}} \left[Y_{\sigma \vee \zeta} + \int_\zeta^{\sigma \vee \zeta} f(s, \theta_s^{Q_n^{(k)}}) ds \middle| \mathcal{F}_\nu \right] &= E_{Q_n^{(k)}} \left[\mathbf{1}_{\{\zeta < \sigma \leq \nu\}} \left(Y_\sigma + \int_\zeta^\sigma f(s, \theta_s^{Q_n^{(k)}}) ds \right) \middle| \mathcal{F}_\nu \right] \\ &= E_{Q_n^{(k)}} \left[\mathbf{1}_{\{\zeta < \sigma \leq \nu\}} Y_{\sigma \wedge \nu} \middle| \mathcal{F}_\nu \right] = \mathbf{1}_{\{\zeta < \sigma \leq \nu\}} Y_{\sigma \wedge \nu} = \mathbf{1}_{\{\zeta < \sigma \leq \nu\}} Y_\sigma; \end{aligned}$$

recall the definitions of the classes $\mathcal{P}_\nu$, $\mathcal{Q}_\nu$ from subsection 1.1. Therefore, (5.43) reduces to

$$\Psi^\sigma(\zeta) \leq E \left[\mathbf{1}_{\{\sigma \leq \nu\}} Y_\sigma + \mathbf{1}_{\{\sigma > \nu\}} E_{Q_n^{(k)}} \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^{Q_n^{(k)}}) ds \middle| \mathcal{F}_\nu \right] \middle| \mathcal{F}_\zeta \right], \quad P\text{-a.s.}$$

We obtain then from (5.38), (5.39) and the Bounded Convergence Theorem, that

$$\begin{aligned} \Psi^\sigma(\zeta) &\leq \lim_{n \rightarrow \infty} \downarrow E \left[\mathbf{1}_{\{\sigma \leq \nu\}} Y_\sigma + \mathbf{1}_{\{\sigma > \nu\}} E_{Q_n^{(k)}} \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^{Q_n^{(k)}}) ds \middle| \mathcal{F}_\nu \right] \middle| \mathcal{F}_\zeta \right] \\ &= E \left[\mathbf{1}_{\{\sigma \leq \nu\}} Y_\sigma + \mathbf{1}_{\{\sigma > \nu\}} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^{Q_n^{(k)}}) ds \middle| \mathcal{F}_\nu \right] \middle| \mathcal{F}_\zeta \right], \quad P\text{-a.s.} \end{aligned}$$

On the other hand, we can deduce from (5.37), (5.40) and the Bounded Convergence Theorem once again that

$$\begin{aligned} \Psi^\sigma(\zeta) &\leq \lim_{k \rightarrow \infty} \downarrow E \left[\mathbf{1}_{\{\sigma \leq \nu\}} Y_\sigma + \mathbf{1}_{\{\sigma > \nu\}} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu^k} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^{Q_n^{(k)}}) ds \middle| \mathcal{F}_\nu \right] \middle| \mathcal{F}_\zeta \right] \\ &= E \left[\mathbf{1}_{\{\sigma \leq \nu\}} Y_\sigma + \mathbf{1}_{\{\sigma > \nu\}} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_{\sigma \vee \nu} + \int_\nu^{\sigma \vee \nu} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \middle| \mathcal{F}_\zeta \right] = E \left[\Psi^\sigma(\nu) \middle| \mathcal{F}_\zeta \right], \quad P\text{-a.s.}, \end{aligned} \quad (5.44)$$

which implies that $\{\Psi^\sigma(t)\}_{t \in [0,T]}$ is a submartingale. Therefore Karatzas and Shreve [1991, Proposition 1.3.14] shows that

$$P \left(\text{the limit } \Psi_t^{\sigma,+} \triangleq \lim_{n \rightarrow \infty} \Psi^\sigma(q_n(t)) \text{ exists for any } t \in [0, T] \right) = 1 \quad (5.45)$$

(where $q_n(t) \triangleq \frac{[2^n t]}{2^n} \wedge T$), and that $\Psi^{\sigma,+}$ is an RCLL process.

Step 3: For any $\nu \in \mathcal{S}_{0,T}$ and $n \in \mathbb{N}$, $q_n(\nu)$ takes values in a finite set $\mathcal{D}_T^n \triangleq ([0, T] \cap \{k2^{-n}\}_{k \in \mathbb{Z}}) \cup \{T\}$. Given an $\lambda \in \mathcal{D}_T^n$, it holds for any $m \geq n$ that $q_m(\lambda) = \lambda$ since $\mathcal{D}_T^n \subset \mathcal{D}_T^m$. It follows from (5.45) that

$$\Psi_\lambda^{\sigma,+} = \lim_{m \rightarrow \infty} \Psi^\sigma(q_m(\lambda)) = \Psi^\sigma(\lambda), \quad P\text{-a.s.}$$

Then one can deduce from (5.42) that

$$\Psi_{q_n(\nu)}^{\sigma,+} = \sum_{\lambda \in \mathcal{D}_T^n} \mathbf{1}_{\{q_n(\nu)=\lambda\}} \Psi_{\lambda}^{\sigma,+} = \sum_{\lambda \in \mathcal{D}_T^n} \mathbf{1}_{\{q_n(\nu)=\lambda\}} \Psi^{\sigma}(\lambda) = \sum_{\lambda \in \mathcal{D}_T^n} \mathbf{1}_{\{q_n(\nu)=\lambda\}} \Psi^{\sigma}(q_n(\nu)) = \Psi^{\sigma}(q_n(\nu)), \quad P\text{-a.s.}$$

Thus the right-continuity of the process $\Psi^{\sigma,+}$ implies that

$$\Psi_{\nu}^{\sigma,+} = \lim_{n \rightarrow \infty} \Psi_{q_n(\nu)}^{\sigma,+} = \lim_{n \rightarrow \infty} \Psi^{\sigma}(q_n(\nu)), \quad P\text{-a.s.} \quad (5.46)$$

Hence (5.44), (5.41) and the Bounded Convergence Theorem imply

$$\Psi^{\sigma}(\nu) \leq \lim_{n \rightarrow \infty} E[\Psi^{\sigma}(q_n(\nu)) | \mathcal{F}_{\nu}] = E[\Psi_{\nu}^{\sigma,+} | \mathcal{F}_{\nu}] = \Psi_{\nu}^{\sigma,+}, \quad P\text{-a.s.} \quad (5.47)$$

In the last equality we used the fact that $\Psi_{\nu}^{\sigma,+} = \lim_{n \rightarrow \infty} \Psi^{\sigma}(q_n(\nu)) \in \mathcal{F}_{\nu}$, thanks to the right-continuity of the Brownian filtration $\mathbf{F}$.

Step 4: Set $\nu, \gamma \in \mathcal{S}_{0,T}$ and $\zeta \triangleq \tau(\nu) \wedge \gamma$, $\zeta_n \triangleq \tau(\nu) \wedge q_n(\gamma)$, $\forall n \in \mathbb{N}$. Now, let $\sigma \in \mathcal{S}_{\zeta,T}$. Since $\lim_{n \rightarrow \infty} \uparrow \mathbf{1}_{\{\tau(\nu) > q_n(\gamma)\}} = \mathbf{1}_{\{\tau(\nu) > \gamma\}}$ and

$$\{\tau(\nu) > \gamma\} \subset \{q_n(\gamma) = q_n(\tau(\nu) \wedge \gamma)\}, \quad \{\tau(\nu) > q_n(\gamma)\} \subset \{q_n(\gamma) = \tau(\nu) \wedge q_n(\gamma)\}, \quad \forall n \in \mathbb{N},$$

one can deduce from (5.47), (5.46), and (5.42) that

$$\begin{aligned} \mathbf{1}_{\{\tau(\nu) > \gamma\}} \Psi^{\sigma}(\zeta) &\leq \mathbf{1}_{\{\tau(\nu) > \gamma\}} \Psi_{\zeta}^{\sigma,+} = \mathbf{1}_{\{\tau(\nu) > \gamma\}} \lim_{n \rightarrow \infty} \Psi^{\sigma}(q_n(\zeta)) = \lim_{n \rightarrow \infty} \mathbf{1}_{\{\tau(\nu) > \gamma\}} \Psi^{\sigma}(q_n(\tau(\nu) \wedge \gamma)) \\ &= \lim_{n \rightarrow \infty} \mathbf{1}_{\{\tau(\nu) > \gamma\}} \Psi^{\sigma}(q_n(\gamma)) = \lim_{n \rightarrow \infty} \mathbf{1}_{\{\tau(\nu) > q_n(\gamma)\}} \Psi^{\sigma}(q_n(\gamma)) \\ &= \lim_{n \rightarrow \infty} \mathbf{1}_{\{\tau(\nu) > q_n(\gamma)\}} \Psi^{\sigma}(\tau(\nu) \wedge q_n(\gamma)) = \mathbf{1}_{\{\tau(\nu) > \gamma\}} \lim_{n \rightarrow \infty} \Psi^{\sigma}(\zeta_n), \quad P\text{-a.s.} \end{aligned} \quad (5.48)$$

For any $n \in \mathbb{N}$, we see from (3.14) and Lemma 2.1 that

$$\begin{aligned} V(\zeta_n) &= \underline{V}(\zeta_n) = \operatorname{ess\,sup}_{\beta \in \mathcal{S}_{\zeta_n,T}} \left(\operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\zeta_n}} E_Q \left[Y_{\beta} + \int_{\zeta_n}^{\beta} f(s, \theta_s^Q) ds \middle| \mathcal{F}_{\zeta_n} \right] \right) \\ &\geq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\zeta_n}} E_Q \left[Y_{\sigma \vee \zeta_n} + \int_{\zeta_n}^{\sigma \vee \zeta_n} f(s, \theta_s^Q) ds \middle| \mathcal{F}_{\zeta_n} \right] \\ &= \operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\zeta_n}} E_Q \left[\mathbf{1}_{\{\sigma \leq \zeta_n\}} Y_{\zeta_n} + \mathbf{1}_{\{\sigma > \zeta_n\}} \left(Y_{\sigma \vee \zeta_n} + \int_{\zeta_n}^{\sigma \vee \zeta_n} f(s, \theta_s^Q) ds \right) \middle| \mathcal{F}_{\zeta_n} \right] \\ &= \operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\zeta_n}} \left(\mathbf{1}_{\{\sigma \leq \zeta_n\}} Y_{\zeta_n} + \mathbf{1}_{\{\sigma > \zeta_n\}} E_Q \left[Y_{\sigma \vee \zeta_n} + \int_{\zeta_n}^{\sigma \vee \zeta_n} f(s, \theta_s^Q) ds \middle| \mathcal{F}_{\zeta_n} \right] \right) \\ &= \mathbf{1}_{\{\sigma \leq \zeta_n\}} Y_{\zeta_n} + \mathbf{1}_{\{\sigma > \zeta_n\}} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\zeta_n}} E_Q \left[Y_{\sigma \vee \zeta_n} + \int_{\zeta_n}^{\sigma \vee \zeta_n} f(s, \theta_s^Q) ds \middle| \mathcal{F}_{\zeta_n} \right], \quad P\text{-a.s.} \end{aligned}$$

Since $\{\tau(\nu) \leq \gamma\} \subset \{\zeta_n = \zeta = \tau(\nu)\}$ and $\{\sigma > \zeta_n\} \subset \{\sigma > \zeta\}$, it follows from (3.16) that

$$\begin{aligned} V(\zeta_n) &\geq \mathbf{1}_{\{\sigma \leq \zeta_n\}} Y_{\zeta_n} + \mathbf{1}_{\{\sigma > \zeta_n, \tau(\nu) > \gamma\}} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\zeta_n}} E_Q \left[Y_{\sigma \vee \zeta_n} + \int_{\zeta_n}^{\sigma \vee \zeta_n} f(s, \theta_s^Q) ds \middle| \mathcal{F}_{\zeta_n} \right] \\ &\quad + \mathbf{1}_{\{\sigma > \zeta_n, \tau(\nu) \leq \gamma\}} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\zeta}} E_Q \left[Y_{\sigma \vee \zeta} + \int_{\zeta}^{\sigma \vee \zeta} f(s, \theta_s^Q) ds \middle| \mathcal{F}_{\zeta} \right] \\ &= \mathbf{1}_{\{\sigma \leq \zeta_n\}} Y_{\zeta_n} + \mathbf{1}_{\{\sigma > \zeta_n, \tau(\nu) > \gamma\}} \Psi^{\sigma}(\zeta_n) + \mathbf{1}_{\{\sigma > \zeta_n, \tau(\nu) \leq \gamma\}} \Psi^{\sigma}(\zeta), \quad P\text{-a.s.} \end{aligned}$$

As $n \rightarrow \infty$, the right-continuity of processes Y , (5.48) as well as Lemma 2.1 show that

$$\begin{aligned}
\lim_{n \rightarrow \infty} V(\zeta_n) &\geq \mathbf{1}_{\{\sigma=\zeta\}} Y_\zeta + \mathbf{1}_{\{\sigma>\zeta, \tau(\nu)>\gamma\}} \lim_{n \rightarrow \infty} \Psi^\sigma(\zeta_n) + \mathbf{1}_{\{\sigma>\zeta, \tau(\nu)\leq\gamma\}} \Psi^\sigma(\zeta) \\
&\geq \mathbf{1}_{\{\sigma=\zeta\}} Y_\zeta + \mathbf{1}_{\{\sigma>\zeta\}} \Psi^\sigma(\zeta) = \mathbf{1}_{\{\sigma=\zeta\}} Y_\zeta + \mathbf{1}_{\{\sigma>\zeta\}} \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\zeta} E_Q \left[Y_{\sigma \vee \zeta} + \int_\zeta^{\sigma \vee \zeta} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\zeta \right] \\
&= \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\zeta} \left(\mathbf{1}_{\{\sigma=\zeta\}} Y_\zeta + \mathbf{1}_{\{\sigma>\zeta\}} E_Q \left[Y_{\sigma \vee \zeta} + \int_\zeta^{\sigma \vee \zeta} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\zeta \right] \right) \\
&= \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\zeta} E_Q \left[\mathbf{1}_{\{\sigma=\zeta\}} Y_\zeta + \mathbf{1}_{\{\sigma>\zeta\}} \left(Y_\sigma + \int_\zeta^\sigma f(s, \theta_s^Q) ds \right) \middle| \mathcal{F}_\zeta \right] \\
&= \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\zeta} E_Q \left[Y_\sigma + \int_\zeta^\sigma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\zeta \right], \quad P\text{-a.s.}
\end{aligned}$$

Taking the essential supremum of the right-hand-side over $\sigma \in \mathcal{S}_{\zeta, T}$, we obtain

$$\lim_{n \rightarrow \infty} V(\zeta_n) \geq \operatorname{ess\,sup}_{\sigma \in \mathcal{S}_{\zeta, T}} \left(\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\zeta} E_Q \left[Y_\sigma + \int_\zeta^\sigma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\zeta \right] \right) = \underline{V}(\zeta) = V(\zeta), \quad P\text{-a.s.} \quad (5.49)$$

Let us show the reverse inequality. Fix $Q \in \mathcal{Q}_\zeta$ and $n \in \mathbb{N}$. For any $k, m \in \mathbb{N}$, the predictable process

$$\theta_t^{Q_n^{m,k}} \triangleq \mathbf{1}_{\{\zeta_n < t \leq \delta_m^{Q,n}\}} \mathbf{1}_{A_{\zeta,k}^Q} \theta_t^Q, \quad t \in [0, T]$$

induces a probability measure $Q_n^{m,k} \in \mathcal{Q}_{\zeta_n}^k$ by $dQ_n^{m,k} \triangleq \mathcal{E} \left(\theta^{Q_n^{m,k}} \bullet B \right)_T dP$, where $\delta_m^{Q,n}$ is defined by

$$\delta_m^{Q,n} \triangleq \inf \left\{ t \in [\zeta_n, T] : \int_{\zeta_n}^t f(s, \theta_s^Q) ds > m \right\} \wedge T, \quad m \in \mathbb{N}.$$

For any $\beta \in \mathcal{S}_{\zeta_n, T}$, using arguments similar to those that lead to (5.7), we obtain

$$\begin{aligned}
E_{Q_n^{m,k}} \left[Y_\beta + \int_{\zeta_n}^\beta f(s, \theta_s^{Q_n^{m,k}}) ds \middle| \mathcal{F}_{\zeta_n} \right] &\leq (\|Y\|_\infty + m) \cdot E \left[\left| Z_{\zeta_n, T}^{Q_n^{m,k}} - Z_{\zeta_n, \delta_m^{Q,n}}^Q \right| \middle| \mathcal{F}_{\zeta_n} \right] \\
&\quad + \|Y\|_\infty \cdot E \left[\left| Z_{\zeta_n, \delta_m^{Q,n}}^Q - Z_{\zeta_n, T}^Q \right| \middle| \mathcal{F}_{\zeta_n} \right] + E_Q \left[Y_\beta + \int_{\zeta_n}^\beta f(s, \theta_s^Q) ds \middle| \mathcal{F}_{\zeta_n} \right], \quad P\text{-a.s.}
\end{aligned}$$

Then taking the essential supremum of both sides over $\beta \in \mathcal{S}_{\zeta_n, T}$ yields that

$$\begin{aligned}
\operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\zeta_n}^k} R^Q(\zeta_n) &\leq R^{Q_n^{m,k}}(\zeta_n) \leq (\|Y\|_\infty + m) E \left[\left| Z_{\zeta_n, T}^{Q_n^{m,k}} - Z_{\zeta_n, \delta_m^{Q,n}}^Q \right| \middle| \mathcal{F}_{\zeta_n} \right] \\
&\quad + \|Y\|_\infty \cdot E \left[\left| Z_{\zeta_n, \delta_m^{Q,n}}^Q - Z_{\zeta_n, T}^Q \right| \middle| \mathcal{F}_{\zeta_n} \right] + R^Q(\zeta_n), \quad P\text{-a.s.} \quad (5.50)
\end{aligned}$$

Just as in (5.8), we can show that

$$\lim_{k \rightarrow \infty} E \left[\left| Z_{\zeta_n, T}^{Q_n^{m,k}} - Z_{\zeta_n, \delta_m^{Q,n}}^Q \right| \middle| \mathcal{F}_{\zeta_n} \right] = 0, \quad P\text{-a.s.}$$

Therefore, letting $k \rightarrow \infty$ in (5.50), we know from Lemma 3.1 (2) that

$$V(\zeta_n) = \lim_{k \rightarrow \infty} \downarrow \operatorname{ess\,inf}_{Q \in \mathcal{Q}_{\zeta_n}^k} R^Q(\zeta_n) \leq \|Y\|_\infty \cdot E \left[\left| Z_{\zeta_n, \delta_m^{Q,n}}^Q - Z_{\zeta_n, T}^Q \right| \middle| \mathcal{F}_{\zeta_n} \right] + R^Q(\zeta_n), \quad P\text{-a.s.} \quad (5.51)$$

Next, by analogy with (5.10), we have $\lim_{m \rightarrow \infty} E \left(\left| Z_{\zeta_n, \delta_m^{Q,n}}^Q - Z_{\zeta_n, T}^Q \right| \middle| \mathcal{F}_{\zeta_n} \right) = 0$, P -a.s. Letting $m \rightarrow \infty$ in (5.51), we obtain $V(\zeta_n) \leq R^Q(\zeta_n) = R_{\zeta_n}^{Q,0}$, P -a.s. from (3.3). Then the right-continuity of the process $R^{Q,0}$, as well as (3.3), imply that

$$\overline{\lim}_{n \rightarrow \infty} V(\zeta_n) \leq \lim_{n \rightarrow \infty} R_{\zeta_n}^{Q,0} = R_\zeta^{Q,0} = R^Q(\zeta), \quad P\text{-a.s.}$$

Taking the essential infimum of $R^Q(\zeta)$ over $Q \in \mathcal{Q}_\zeta$ yields $\overline{\lim}_{n \rightarrow \infty} V(\zeta_n) \leq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\zeta} R^Q(\zeta) = \overline{V}(\zeta) = V(\zeta)$, P -a.s. This inequality, together with (5.49), shows that

$$\lim_{n \rightarrow \infty} V(\tau(\nu) \wedge q_n(\gamma)) = V(\tau(\nu) \wedge \gamma), \quad P\text{-a.s.} \quad (5.52)$$

Step 5: Now fix $\nu \in \mathcal{S}_{0,T}$. It is clear that $P \in \mathcal{Q}_\nu$ and that $\theta^P \equiv 0$. For any $t \in [0, T]$, (3.17) implies that

$$\mathbf{1}_{\{t \geq \nu\}} V(\tau(\nu) \wedge t) = \mathbf{1}_{\{t \geq \nu\}} V(\tau(\nu) \wedge (t \vee \nu)), \quad P\text{-a.s.},$$

since $\{t \geq \nu\} \subset \{\tau(\nu) \wedge t = \tau(\nu) \wedge (t \vee \nu)\}$. Then we can deduce from (3.15), (f3), and (3.14) that for any $s \in [0, t]$

$$\begin{aligned} \mathbf{1}_{\{s \geq \nu\}} V(\tau(\nu) \wedge s) &= \mathbf{1}_{\{s \geq \nu\}} V(\tau(\nu) \wedge (s \vee \nu)) \\ &\leq \mathbf{1}_{\{s \geq \nu\}} E \left[V(\tau(\nu) \wedge (t \vee \nu)) + \int_{\tau(\nu) \wedge (s \vee \nu)}^{\tau(\nu) \wedge (t \vee \nu)} f(r, \theta_r^P) dr \middle| \mathcal{F}_{\tau(\nu) \wedge (s \vee \nu)} \right] \\ &= \mathbf{1}_{\{s \geq \nu\}} E \left[V(\tau(\nu) \wedge (t \vee \nu)) \middle| \mathcal{F}_{\tau(\nu) \wedge s} \right] = E \left[\mathbf{1}_{\{s \geq \nu\}} V(\tau(\nu) \wedge (t \vee \nu)) \middle| \mathcal{F}_{\tau(\nu) \wedge s} \right] \\ &\leq E \left[\mathbf{1}_{\{t \geq \nu\}} V(\tau(\nu) \wedge (t \vee \nu)) + \mathbf{1}_{\{t \geq \nu > s\}} \|Y\|_\infty \middle| \mathcal{F}_{\tau(\nu) \wedge s} \right] \\ &= E \left[E \left[\mathbf{1}_{\{t \geq \nu\}} (V(\tau(\nu) \wedge t) + \|Y\|_\infty) \middle| \mathcal{F}_{\tau(\nu)} \right] \middle| \mathcal{F}_s \right] - \mathbf{1}_{\{s \geq \nu\}} \|Y\|_\infty \\ &= E \left[\mathbf{1}_{\{t \geq \nu\}} (V(\tau(\nu) \wedge t) + \|Y\|_\infty) \middle| \mathcal{F}_s \right] - \mathbf{1}_{\{s \geq \nu\}} \|Y\|_\infty, \quad P\text{-a.s.}, \end{aligned}$$

which shows that $\left\{ \mathbf{1}_{\{t \geq \nu\}} (V(\tau(\nu) \wedge t) + \|Y\|_\infty) \right\}_{t \in [0, T]}$ is a submartingale. Hence it follows from Karatzas and Shreve [1991, Proposition 1.3.14] that

$$P \left(\text{the limit } V_t^{0, \nu} \triangleq \lim_{n \rightarrow \infty} \mathbf{1}_{\{q_n(t) \geq \nu\}} V(\tau(\nu) \wedge q_n(t)) \text{ exists for any } t \in [0, T] \right) = 1,$$

and that $V^{0, \nu}$ is an RCLL process.

Let $\zeta \in \mathcal{S}_{0,T}^*$ take values in a finite set $\{t_1 < \dots < t_m\}$. For any $\lambda \in \{1 \dots m\}$ and $n \in \mathbb{N}$, since $\{\zeta = t_\lambda\} \subset \{\tau(\nu) \wedge q_n(\zeta) = \tau(\nu) \wedge q_n(t_\lambda)\}$, one can deduce from (3.17) that

$$\mathbf{1}_{\{\zeta = t_\lambda\}} V(\tau(\nu) \wedge q_n(\zeta)) = \mathbf{1}_{\{\zeta = t_\lambda\}} V(\tau(\nu) \wedge q_n(t_\lambda)), \quad P\text{-a.s.}$$

As $n \rightarrow \infty$, (5.52) shows

$$\begin{aligned} \mathbf{1}_{\{\zeta = t_\lambda\}} V_\zeta^{0, \nu} &= \mathbf{1}_{\{\zeta = t_\lambda\}} V_{t_\lambda}^{0, \nu} = \mathbf{1}_{\{t_\lambda \geq \nu\}} \lim_{n \rightarrow \infty} \mathbf{1}_{\{\zeta = t_\lambda\}} V(\tau(\nu) \wedge q_n(t_\lambda)) \\ &= \mathbf{1}_{\{t_\lambda \geq \nu\}} \lim_{n \rightarrow \infty} \mathbf{1}_{\{\zeta = t_\lambda\}} V(\tau(\nu) \wedge q_n(\zeta)) = \mathbf{1}_{\{\zeta \geq \nu\}} \mathbf{1}_{\{\zeta = t_\lambda\}} V(\tau(\nu) \wedge \zeta), \quad P\text{-a.s.} \end{aligned}$$

Summing the above expression over λ , we obtain $V_\zeta^{0, \nu} = \mathbf{1}_{\{\zeta \geq \nu\}} V(\tau(\nu) \wedge \zeta)$, P -a.s. Then for any $\gamma \in \mathcal{S}_{0,T}$, the right-continuity of the process $V^{0, \nu}$ and (5.52) imply

$$V_\gamma^{0, \nu} = \lim_{n \rightarrow \infty} V_{q_n(\gamma)}^{0, \nu} = \lim_{n \rightarrow \infty} \mathbf{1}_{\{q_n(\gamma) \geq \nu\}} V(\tau(\nu) \wedge q_n(\gamma)) = \mathbf{1}_{\{\gamma \geq \nu\}} V(\tau(\nu) \wedge \gamma), \quad P\text{-a.s.},$$

proving (3.18). In particular, $V^{0, \nu}$ is an RCLL modification of the process $\{\mathbf{1}_{\{t \geq \nu\}} V(\tau(\nu) \wedge t)\}_{t \in [0, T]}$.

Proof of Theorem 3.2: Proof of (2). Proposition 3.2 and (3.18) imply that $V_{\tau(\nu)}^{0, \nu} = V(\tau(\nu)) = Y_{\tau(\nu)}$, P -a.s. Hence we can deduce from the right-continuity of processes $V^{0, \nu}$ and Y that $\tau_V(\nu)$ in (3.19) is a stopping time belonging to $\mathcal{S}_{\nu, \tau(\nu)}$ and that

$$Y_{\tau_V(\nu)} = V_{\tau_V(\nu)}^{0, \nu} = V(\tau_V(\nu)), \quad P\text{-a.s.},$$

where the second equality is due to (3.18). Then it follows from (3.15) that for any $Q \in \mathcal{Q}_\nu$

$$V(\nu) \leq E_Q \left[V(\tau_V(\nu)) + \int_\nu^{\tau_V(\nu)} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] = E_Q \left[Y_{\tau_V(\nu)} + \int_\nu^{\tau_V(\nu)} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.}$$

Taking the essential infimum of the right-hand-side over $Q \in \mathcal{Q}_\nu$ yields that

$$\begin{aligned} V(\nu) &\leq \operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_{\tau_V(\nu)} + \int_\nu^{\tau_V(\nu)} f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \\ &\leq \operatorname{ess\,sup}_{\gamma \in \mathcal{S}_{\nu, T}} \left(\operatorname{ess\,inf}_{Q \in \mathcal{Q}_\nu} E_Q \left[Y_\gamma + \int_\nu^\gamma f(s, \theta_s^Q) ds \middle| \mathcal{F}_\nu \right] \right) = \underline{V}(\nu) = V(\nu), \quad P\text{-a.s.}, \end{aligned}$$

from which the claim follows. $\square$

5.2 Proof of the Results in Section 4

Proof of Theorem 4.1: It is easy to see from (i) that

$$Y_{\sigma_*} = V(\sigma_*) = R^{Q^*}(\sigma_*), \quad P\text{-a.s.} \quad (5.53)$$

which together with (ii) and (iii) shows that for any $Q \in \mathcal{Q}_0$

$$E_{Q^*} [Y_{\sigma_*}^{Q^*}] = E_{Q^*} [V^{Q^*}(\sigma_*)] = V^{Q^*}(0) = V(0) \leq E_Q [V^Q(\sigma_*)] = E_Q [Y_{\sigma_*}^Q].$$

Thus the second inequality in (4.1) holds for (Q^*, σ_*) . Now we show that (Q^*, σ_*) satisfies the first inequality in (4.1) in three steps:

- When $\nu \in \mathcal{S}_{0, \sigma_*}$, property (iii) and (5.53) imply that

$$Y_\nu^{Q^*} \leq V^{Q^*}(\nu) = E_{Q^*} [V^{Q^*}(\sigma_*) | \mathcal{F}_\nu] = E_{Q^*} [Y_{\sigma_*}^{Q^*} | \mathcal{F}_\nu], \quad P\text{-a.s.} \quad (5.54)$$

Taking the expectation E_{Q^*} on both sides yields that $E_{Q^*} [Y_\nu^{Q^*}] \leq E_{Q^*} [Y_{\sigma_*}^{Q^*}]$.

- When $\nu \in \mathcal{S}_{\sigma_*, T}$, it follows from (5.53) that

$$\begin{aligned} E_{Q^*} [Y_\nu^{Q^*}] &= E_{Q^*} \left[E_{Q^*} \left[Y_\nu + \int_{\sigma_*}^\nu f(s, \theta_s^{Q^*}) ds \middle| \mathcal{F}_{\sigma_*} \right] + \int_0^{\sigma_*} f(s, \theta_s^{Q^*}) ds \right] \\ &\leq E_{Q^*} \left[R^{Q^*}(\sigma_*) + \int_0^{\sigma_*} f(s, \theta_s^{Q^*}) ds \right] = E_{Q^*} [Y_{\sigma_*}^{Q^*}]. \end{aligned}$$

- For a general stopping time $\nu \in \mathcal{S}_{0, T}$, let us define $\nu_1 = \nu \wedge \sigma_* \in \mathcal{S}_{0, \sigma_*}$ and $\nu_2 = \nu \vee \sigma_* \in \mathcal{S}_{\sigma_*, T}$. Since $\{\nu \leq \sigma_*\} \in \mathcal{F}_{\nu \wedge \sigma_*} = \mathcal{F}_{\nu_1}$, one can deduce from (5.54) that

$$\begin{aligned} E_{Q^*} [Y_\nu^{Q^*}] &= E_{Q^*} \left[E_{Q^*} \left[\mathbf{1}_{\{\nu \leq \sigma_*\}} Y_{\nu_1}^{Q^*} + \mathbf{1}_{\{\nu > \sigma_*\}} Y_{\nu_2}^{Q^*} \middle| \mathcal{F}_{\sigma_*} \right] \right] = E_{Q^*} \left[\mathbf{1}_{\{\nu \leq \sigma_*\}} Y_{\nu_1}^{Q^*} + \mathbf{1}_{\{\nu > \sigma_*\}} E_{Q^*} [Y_{\nu_2}^{Q^*} | \mathcal{F}_{\sigma_*}] \right] \\ &\leq E_{Q^*} \left[\mathbf{1}_{\{\nu \leq \sigma_*\}} Y_{\nu_1}^{Q^*} + \mathbf{1}_{\{\nu > \sigma_*\}} \left(R^{Q^*}(\sigma_*) + \int_0^{\sigma_*} f(s, \theta_s^{Q^*}) ds \right) \right] \\ &= E_{Q^*} \left[\mathbf{1}_{\{\nu \leq \sigma_*\}} Y_{\nu_1}^{Q^*} + \mathbf{1}_{\{\nu > \sigma_*\}} Y_{\sigma_*}^{Q^*} \right] = E_{Q^*} \left[\mathbf{1}_{\{\nu \leq \sigma_*\}} Y_{\nu_1}^{Q^*} + \mathbf{1}_{\{\nu > \sigma_*\}} E_{Q^*} [Y_{\sigma_*}^{Q^*} | \mathcal{F}_{\nu_1}] \right] \\ &\leq E_{Q^*} \left[\mathbf{1}_{\{\nu \leq \sigma_*\}} E_{Q^*} [Y_{\sigma_*}^{Q^*} | \mathcal{F}_{\nu_1}] + \mathbf{1}_{\{\nu > \sigma_*\}} E_{Q^*} [Y_{\sigma_*}^{Q^*} | \mathcal{F}_{\nu_1}] \right] = E_{Q^*} [Y_{\sigma_*}^{Q^*}]. \end{aligned} \quad \square$$

Proof of Lemma 4.1: Fix $t \in [0, T]$. For any $\gamma \in \mathcal{S}_{\nu \vee t, T}$, we see from (4.5) that

$$\tilde{\Gamma}_{\nu \vee t} = \tilde{\Gamma}_\gamma + \int_{\nu \vee t}^\gamma f(s, \theta_s^{*, \nu}) ds + \tilde{K}_\gamma - \tilde{K}_{\nu \vee t} - \int_{\nu \vee t}^\gamma \tilde{Z}_s dB_s^{Q^{*, \nu}}, \quad P\text{-a.s.}$$

Applying $E_{Q^*,\nu}[\cdot|\mathcal{F}_{\nu\vee t}]$ to both sides, we obtain

$$\tilde{\Gamma}_{\nu\vee t} = E_{Q^*,\nu} \left[\tilde{\Gamma}_\gamma + \int_{\nu\vee t}^\gamma f(s, \theta_s^{*,\nu}) ds + \tilde{K}_\gamma - \tilde{K}_{\nu\vee t} \middle| \mathcal{F}_{\nu\vee t} \right] \quad (5.55)$$

$$\geq E_{Q^*,\nu} \left[Y_\gamma + \int_{\nu\vee t}^\gamma f(s, \theta_s^{*,\nu}) ds \middle| \mathcal{F}_{\nu\vee t} \right], \quad P\text{-a.s.} \quad (5.56)$$

Let $\sigma_{\nu\vee t}^* \triangleq \inf\{s \in [\nu\vee t, T] : \tilde{\Gamma}_s = Y_s\} \in \mathcal{S}_{\nu\vee t, T}$. The flat-off condition satisfied by $(\tilde{\Gamma}, \tilde{Z}, \tilde{K})$, and the continuity of $\tilde{K}$, imply that

$$0 = \int_{[\nu\vee t, \sigma_{\nu\vee t}^*)} \mathbf{1}_{\{\tilde{\Gamma}_s > Y_s\}} d\tilde{K}_s = \int_{[\nu\vee t, \sigma_{\nu\vee t}^*)} d\tilde{K}_s = \lim_{s \nearrow \sigma_{\nu\vee t}^*} \tilde{K}_s - \tilde{K}_{\nu\vee t} = \tilde{K}_{\sigma_{\nu\vee t}^*} - \tilde{K}_{\nu\vee t}, \quad P\text{-a.s.}$$

Hence, taking $\gamma = \sigma_{\nu\vee t}^*$ in (5.55), we obtain the P -a.s. property $\tilde{\Gamma}_{\nu\vee t} = E_{Q^*,\nu} \left[Y_{\sigma_{\nu\vee t}^*} + \int_{\nu\vee t}^{\sigma_{\nu\vee t}^*} f(s, \theta_s^{*,\nu}) ds \middle| \mathcal{F}_{\nu\vee t} \right]$ which, together with (5.56) and (3.3), shows that

$$\tilde{\Gamma}_{\nu\vee t} = \operatorname{ess\,sup}_{\gamma \in \mathcal{S}_{\nu\vee t, T}} E_{Q^*,\nu} \left[Y_\gamma + \int_{\nu\vee t}^\gamma f(s, \theta_s^{*,\nu}) ds \middle| \mathcal{F}_{\nu\vee t} \right] = R^{Q^*,\nu}(\nu\vee t) = R_{\nu\vee t}^{Q^*,\nu,0}, \quad P\text{-a.s.}$$

Then the right-continuity of the processes $\tilde{\Gamma}$ and $R^{Q^*,\nu,0}$ implies (4.6). $\square$

Proof of Theorem 4.2 : We shall show that $(Q^*, \tau^{Q^*}(0))$ satisfies conditions (i)-(iii) of Theorem 4.1:

- 1) It follows easily from Proposition 3.1 that $Y_{\tau^{Q^*}(0)} = R_{\tau^{Q^*}(0)}^{Q^*,0} = R^{Q^*}(\tau^{Q^*}(0))$, P -a.s.
- 2) For any $k \in \mathbb{N}$ and $Q \in \mathcal{Q}_0^k$, we can deduce from (4.9), the right-continuity of processes $R^{Q^*,0}$ and $\tilde{\Gamma}$, as well as (4.8) that P -a.s.

$$R_t^{Q^*,0} = \tilde{\Gamma}_t \leq R_t^{Q,0}, \quad \forall t \in [0, T].$$

In particular, we have $Y_{\tau^Q(0)} \leq R_{\tau^Q(0)}^{Q^*,0} = R_{\tau^Q(0)}^{Q,0} = Y_{\tau^Q(0)}$, P -a.s. Hence $Y_{\tau^Q(0)} = R_{\tau^Q(0)}^{Q^*,0}$, P -a.s., which implies further that $\tau^{Q^*}(0) \leq \tau^Q(0)$, P -a.s. Taking the essential infimum of right-hand-side over $Q \in \mathcal{Q}_0^k$ and letting $k \rightarrow \infty$, we deduce that, in the notation of (3.11), we have $\tau^{Q^*}(0) \leq \lim_{k \rightarrow \infty} \downarrow \operatorname{ess\,inf}_{Q \in \mathcal{Q}_0^k} \tau^Q(0) = \tau(0)$, P -a.s. Then (3.15) shows $V(0) \leq E_Q[V^Q(\tau^{Q^*}(0))]$ for any $Q \in \mathcal{Q}_0$.

- 3) For any $\nu \in \mathcal{S}_{0, \tau^{Q^*}(0)}$, and since $\nu \leq \tau^{Q^*}(0) \leq \tau^{Q^*}(\nu)$ holds P -a.s., one can deduce from (4.9) and (3.4) that

$$\begin{aligned} V^{Q^*}(\nu) &= R^{Q^*}(\nu) + \int_0^\nu f(s, \theta_s^*) ds = E_{Q^*} \left[R^{Q^*}(\tau^{Q^*}(0)) + \int_\nu^{\tau^{Q^*}(0)} f(s, \theta_s^*) ds \middle| \mathcal{F}_\nu \right] + \int_0^\nu f(s, \theta_s^*) ds \\ &= E_{Q^*} \left[R^{Q^*}(\tau^{Q^*}(0)) + \int_0^{\tau^{Q^*}(0)} f(s, \theta_s^*) ds \middle| \mathcal{F}_\nu \right] = E_{Q^*} \left[V^{Q^*}(\tau^{Q^*}(0)) \middle| \mathcal{F}_\nu \right], \quad P\text{-a.s.} \end{aligned} \quad \square$$

References

- E. Bayraktar and S. Yao. Optimal stopping for nonlinear expectations. Technical report, University of Michigan, 2009. Available at <http://arxiv.org/abs/0905.3601>.
- V. E. Beneš. Existence of optimal strategies based on specified information, for a class of stochastic decision problems. *SIAM J. Control*, 8:179–188, 1970. ISSN 0363-0129.
- J. Bion-Nadal. Time consistent dynamic risk processes. *Stochastic Process. Appl.*, 119(2):633–654, 2009. ISSN 0304-4149.

- P. Cheridito, F. Delbaen, and M. Kupper. Dynamic monetary risk measures for bounded discrete-time processes. *Electron. J. Probab.*, 11:no. 3, 57–106 (electronic), 2006. ISSN 1083-6489.
- F. Delbaen. The structure of m -stable sets and in particular of the set of risk neutral measures. In *In memoriam Paul-André Meyer: Séminaire de Probabilités XXXIX*, volume 1874 of *Lecture Notes in Math.*, pages 215–258. Springer, Berlin, 2006.
- F. Delbaen, S. Peng, and E. Rosazza-Gianin. Representation of the penalty term of dynamic concave utilities. Technical report, ETH, 2009. Available at <http://arxiv.org/abs/0802.1121>.
- N. El Karoui. Les aspects probabilistes du contrôle stochastique. In *Ninth Saint Flour Probability Summer School—1979 (Saint Flour, 1979)*, volume 876 of *Lecture Notes in Math.*, pages 73–238. Springer, Berlin, 1981.
- N. El Karoui, C. Kapoudjian, E. Pardoux, S. Peng, and M. C. Quenez. Reflected solutions of backward SDE's, and related obstacle problems for PDE's. *Ann. Probab.*, 25(2):702–737, 1997. ISSN 0091-1798.
- R. J. Elliott. *Stochastic calculus and applications*, volume 18 of *Applications of Mathematics (New York)*. Springer-Verlag, New York, 1982. ISBN 0-387-90763-7.
- H. Föllmer and I. Penner. Convex risk measures and the dynamics of their penalty functions. *Statist. Decisions*, 24(1):61–96, 2006. ISSN 0721-2631.
- H. Föllmer and A. Schied. *Stochastic finance*, volume 27 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, extended edition, 2004. ISBN 3-11-018346-3. An introduction in discrete time.
- I. Karatzas and S. E. Shreve. *Brownian motion and stochastic calculus*, volume 113 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1991. ISBN 0-387-97655-8.
- I. Karatzas and S. E. Shreve. *Methods of mathematical finance*, volume 39 of *Applications of Mathematics (New York)*. Springer-Verlag, New York, 1998. ISBN 0-387-94839-2.
- I. Karatzas and I. M. Zamfirescu. Martingale approach to stochastic control with discretionary stopping. *Appl. Math. Optim.*, 53(2):163–184, 2006. ISSN 0095-4616.
- I. Karatzas and I. M. Zamfirescu. Martingale approach to stochastic differential games of control and stopping. *Ann. Probab.*, 36(4):1495–1527, 2008. ISSN 0091-1798.
- N. Kazamaki. *Continuous exponential martingales and BMO*, volume 1579 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1994. ISBN 3-540-58042-5.
- S. Klöppel and M. Schweizer. Dynamic indifference valuation via convex risk measures. *Math. Finance*, 17(4):599–627, 2007. ISSN 0960-1627.
- M. Kobylanski, J. P. Lepeltier, M. C. Quenez, and S. Torres. Reflected BSDE with superlinear quadratic coefficient. *Probab. Math. Statist.*, 22(1, Acta Univ. Wratislav. No. 2409):51–83, 2002. ISSN 0208-4147.
- J.-P. Lepeltier. On a general zero-sum stochastic game with stopping strategy for one player and continuous strategy for the other. *Probab. Math. Statist.*, 6(1):43–50, 1985. ISSN 0208-4147.
- J. Neveu. *Discrete-parameter martingales*. North-Holland Publishing Co., Amsterdam, revised edition, 1975. Translated from the French by T. P. Speed, North-Holland Mathematical Library, Vol. 10.
- R. T. Rockafellar. *Convex analysis*. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 1997. ISBN 0-691-01586-4. Reprint of the 1970 original, Princeton Paperbacks.
- D. Williams. *Probability with martingales*. Cambridge Mathematical Textbooks. Cambridge University Press, Cambridge, 1991. ISBN 0-521-40455-X; 0-521-40605-6.